

Durham Research Online

Deposited in DRO:

17 May 2017

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

White, T.S. and Bridgland, D.R. and Limondin-Lozouet, N. and Schreve, D.C. (2017) 'Fossils from Quaternary fluvial archives : sources of biostratigraphical, biogeographical and palaeoclimatic evidence.', Quaternary science reviews., 166 . pp. 150-176.

Further information on publisher's website:

<https://doi.org/10.1016/j.quascirev.2017.04.016>

Publisher's copyright statement:

© 2017 This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Manuscript Number: JQSR-D-16-00468R1

Title: Fossils from Quaternary fluvial archives: sources of biostratigraphical, biogeographical and palaeoclimatic evidence

Article Type: SI:Fluvial Archives Group

Keywords: Fluvial archives, biostratigraphy, palaeoenvironments, biogeography, vertebrates, invertebrates

Corresponding Author: Dr. Tom S White, Ph.D.

Corresponding Author's Institution: University of Cambridge

First Author: Tom S White, Ph.D.

Order of Authors: Tom S White, Ph.D.; David R Bridgland, PhD; Nicole Limondin-Lozouet, PhD; Schreve C Danielle, PhD

Abstract: Fluvial sedimentary archives have the potential to preserve a wide variety of palaeontological evidence, ranging from robust bones and teeth found in coarse gravel aggradations to delicate insect remains and plant macrofossils from fine-grained deposits. Over the last decade, advances in Quaternary biostratigraphy based on vertebrate and invertebrate fossils (primarily mammals and molluscs) have been made in many parts of the world, resulting in improved relative chronologies for fluvial sequences. Complementary fossil groups, such as insects, ostracods and plant macrofossils, are also increasingly used in multi-proxy palaeoclimatic and palaeoenvironmental reconstructions, allowing direct comparison of the climates and environments that prevailed at different times across widely separated regions. This paper reviews these topics on a regional basis, with an emphasis on the latest published information, and represents an update to the 2007 review compiled by the FLAG-inspired IGCP 449 biostratigraphy subgroup. Disparities in the level of detail available for different regions can largely be attributed to varying potential for preservation of fossil material, which is especially poor in areas of non-calcareous bedrock, but to some extent also reflect research priorities in different parts of the world. Recognition of the value of biostratigraphical and palaeoclimatic frameworks, which have been refined over many decades in the 'core regions' for such research (particularly for the late Middle and Late Pleistocene of NW Europe), has focussed attention on the need to accumulate similar palaeontological datasets in areas lacking such long research histories. Although the emerging datasets from these understudied regions currently allow only tentative conclusions to be drawn, they represent an important stage in the development of independent biostratigraphical and palaeoenvironmental schemes, which can then be compared and contrasted.

1 QSR Special Issue: 20 years of Fluvial Archives Group

2
3 **Fossils from Quaternary fluvial archives: sources of**
4 **biostratigraphical, biogeographical and palaeoclimatic evidence**

5
6 Tom S. White^a, David R. Bridgland^b, Nicole Limondin-Lozouet^c, Danielle C.
7 Schreve^d

8
9 ^a Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2 3EJ, UK.

10 ^b Department of Geography, University of Durham, South Road, Durham, DH1 3LE, UK

11 ^c Laboratoire de Géographie Physique, UMR8591 CNRS-Université Paris 1 & UPEC, 1 Place A.
12 Briand, F-92195 Meudon Cedex, France

13 ^d Department of Geography, Royal Holloway University of London, Egham, Surrey, TW20 0EX, UK

14
15 **Abstract**

16 Fluvial sedimentary archives have the potential to preserve a wide variety of
17 palaeontological evidence, ranging from robust bones and teeth found in coarse gravel
18 aggradations to delicate insect remains and plant macrofossils from fine-grained
19 deposits. Over the last decade, advances in Quaternary biostratigraphy based on
20 vertebrate and invertebrate fossils (primarily mammals and molluscs) have been made
21 in many parts of the world, resulting in improved relative chronologies for fluvial
22 sequences. Complementary fossil groups, such as insects, ostracods and plant
23 macrofossils, are also increasingly used in multi-proxy palaeoclimatic and
24 palaeoenvironmental reconstructions, allowing direct comparison of the climates and
25 environments that prevailed at different times across widely separated regions. This
26 paper reviews these topics on a regional basis, with an emphasis on the latest
27 published information, and represents an update to the 2007 review compiled by the
28 FLAG-inspired IGCP 449 biostratigraphy subgroup. Disparities in the level of detail
29 available for different regions can largely be attributed to varying potential for
30 preservation of fossil material, which is especially poor in areas of non-calcareous
31 bedrock, but to some extent also reflect research priorities in different parts of the
32 world. Recognition of the value of biostratigraphical and palaeoclimatic frameworks,
33 which have been refined over many decades in the 'core regions' for such research
34 (particularly for the late Middle and Late Pleistocene of NW Europe), has focussed
35 attention on the need to accumulate similar palaeontological datasets in areas lacking
36 such long research histories. Although the emerging datasets from these understudied
37 regions currently allow only tentative conclusions to be drawn, they represent an
38 important stage in the development of independent biostratigraphical and
39 palaeoenvironmental schemes, which can then be compared and contrasted.

Figures:

Fig. 1 Map showing regions, major river catchments and key fossil sites mentioned in the text.

Fig. 2 Idealized tranverse section through the Lower Thames terrace sequence, east of London, including details of Mammal Assemblage Zones (after Bridgland and Schreve, 2001), updated to include biostratigraphically significant invertebrate species and archaeological data.

Fig. 3 Correlation of Hoxnian fluvial sequences in the Lower Thames with the stratotype at Hoxne (after White et al., 2013). The occurrence of the 'Rhenish' suite of freshwater molluscs is critical to linking the Swanscombe and Clacton sequences and thence to the pollen stratigraphy of the type-Hoxnian (sub-stages labelled). Units for which there is clear palaeontological or sedimentological evidence for the prevailing climate are colour-coded, and tentative correlations with the marine oxygen isotope stages (MIS) of the deep-sea record are also shown.

Fig. 4 Idealized cross-sections through the Middle Trent, Lower Trent and Witham terrace sequences (modified from Bridgland et al., 2015 and White et al., 2017), showing MIS correlation and biostratigraphically significant species.

Fig. 5 Schematic diagrams showing the progressive appearance of forest land snail species during three interglacials in northern France: La Celle (MIS 11), Caours (MIS 5e) and St Germain-le-Vasson (MIS 1). Modified from Limondin-Lozouet and Preece (2014).

Fig. 6 Synthetic curves showing the development of ecological groups of molluscs during MIS 11 at the site of La Celle, northern France, and a comparison with malacological successions from other western European MIS 11 tufa sequences. The appearances of critical species at La Celle that permit correlation with other sequences are highlighted (after Limondin-Lozouet et al., 2015)

Fig. 7 Biostratigraphical scheme for Eastern European mammalian faunas (after Markova, 2007).

Fig. 8 Correlation chart showing faunal changes, the hominin fossil record and significant technological changes in the Indian subcontinent, mainland and insular SE Asia (after Mishra et al., 2010).

Fig. 9 Pleistocene chronology of the Great American Biotic Interchange (GABI), showing details of major faunal exchanges between North and South America that began at around 2.8 Ma (modified from Woodburne, 2010). Marine oxygen isotope stages after Lisiecki and Raymo (2005).

Fig. 10 Biostratigraphical significance of four species of glyptodont (*Neosclerocalyptus*) in the Pleistocene of South America (modified from Zurita et al., 2009a).

Fig. 11 Schematic diagram showing temporal occurrence of megafaunal taxa at Darling Downs, Australia, in relation to the widely-accepted hypothetical megafaunal extinction 'window' and the approximate timing of human arrival on the continent (after Price et al., 2011).

1. Introduction

The study of fossil assemblages recovered from fluvial deposits is well-established as an important element of multi-disciplinary Quaternary research, providing a basis for regional relative dating frameworks (e.g. Gliozzi et al., 1997; Schreve, 2001a; Schreve and Bridgland, 2002; Bridgland et al., 2004; Markova, 2007; Megirian et al., 2010) and for detailed palaeoclimatic and palaeoenvironmental analyses (e.g. Coope, 2010; Schreve and Candy, 2010; Candy et al., 2010, 2015, 2016; Kahlke et al., 2011; Limondin-Lozouet et al., 2010; Rule et al., 2012; White et al., 2013). The contribution of the Fluvial Archives Group (FLAG) to these research areas over the last 20 years has been considerable: two UNESCO-sponsored International Geological Correlation Programme (IGCP) projects, entitled 'Global correlation of Late Cenozoic fluvial deposits' (IGCP 449) and 'Fluvial sequences as evidence for landscape and climatic evolution in the Late Cenozoic' (IGCP 518), ran under the auspices of FLAG between 2000 and 2007 (Bridgland et al., 2007; Westaway et al., 2009). The former included a thematic biostratigraphy subgroup, which compiled data on faunal assemblages from fluvial sequences in different parts of the world, resulting in the publication of a review of progress in faunal correlation of Late Cenozoic fluvial sequences (Schreve et al., 2007); following the conclusion of these IGCP projects, data has continued to accumulate as part of the ongoing efforts of members of FLAG (Cordier et al., 2017).

This paper provides updated reviews, organized on a regional basis, of advances in Pleistocene vertebrate and invertebrate biostratigraphy made during the 20-year life of FLAG; as such, it is intended to be a companion to the report of the IGCP 449 biostratigraphy subgroup (Schreve et al., 2007), which remains a benchmark review of biostratigraphical frameworks derived from fluvial archives around the world. As was the case with that paper, much of the most detailed work has been undertaken in regions such as NW and Central Europe, which have enjoyed long traditions of Pleistocene palaeontological research, particularly for the late Middle and Late Pleistocene (Fig. 1). This review also extends its scope beyond that of its predecessors to include considerations of the palaeoenvironmental, palaeoecological and biogeographical information that can be derived from fossil assemblages, and the potential for Pleistocene fluvial archives to enhance knowledge of long-term Quaternary climate change.

Insert Figure 1 hereabouts

Fluvial archives have several advantages over other types of terrestrial sedimentary sequences that commonly preserve fossils (such as those from lake basins or caves), the most significant of these being the potential chrono-stratigraphical control provided by river terraces, which have great value as regional templates for the terrestrial Quaternary record (Bridgland, 2000, 2006; Antoine et al., 2007; Bridgland and Westaway, 2008a, 2014; Bridgland et al., 2004, 2006, 2017; Mishra et al., 2007; Vandenberghe, 2015). The nature of rivers also means that Pleistocene fluvial sequences have the potential to preserve a variety of plant and animal fossils derived from terrestrial, freshwater and estuarine environments, recording concurrent changes in a diverse range of palaeoenvironmental settings. Lacustrine sequences, although undoubtedly sources of more continuous and higher-resolution longer-timescale records than rivers, usually represent only the deepest part of a lake basin from which the longest sediment sequences can be obtained; such records are therefore often

lacking in fossils indicative of local terrestrial environments (with the notable exception of wind-borne pollen) and are (usually) isolated from marine influences. Conversely, cave sequences, which are often significant repositories of vertebrate fossils (particularly mammals), usually lack pollen and invertebrate assemblages. Fluvial archives therefore have an unrivalled capacity to provide insights into changing Quaternary climates and environments through time, due to the wide range of fossil types available (which include pollen, plant macrofossils, mammals, molluscs, ostracods and insects), coupled with the chronological control provided by river terraces. Together, these proxies can be used to generate local biostratigraphical frameworks, which can then be extended into other regions and into different depositional contexts outside river valleys. Moreover, where fluvial sequences interdigitate with other sediment types, such as glacial or estuarine-marine deposits, biostratigraphical evidence can be used to constrain the age of regional glacial and sea-level histories (e.g. Bridgland et al., 1999; Matoshko et al., 2004; White et al., 2010, 2013, 2017; Vis et al., 2010), which can then provide further stratigraphical ('event stratigraphy') markers. It should be noted, of course, that this potential can only be realized in areas conducive to the preservation of fossils, and that fluvial sequences are subjected to the same taphonomic complications known to affect other Quaternary sedimentary archives.

Much of the recent palaeontological research summarized here has been undertaken as part of multidisciplinary projects, often driven by archaeologically-motivated research questions (see Chauhan et al., 2017). In Europe and Asia, these have often focussed on the timing of hominin occupations relative to changing Pleistocene climate and environments (e.g. Parfitt et al., 2005, 2010; Antoine et al., 2015, 2016; Maddy et al., 2015; Peretto et al., 2015), whereas further afield, in regions such as North America and Australia, attention has been more focussed on potential human impacts, particularly on megafaunal populations (e.g. Prideaux et al., 2010; Prescott et al., 2012; Sandom et al., 2014; Cooper et al., 2015; Stuart, 2015). The potential for fossils to provide important chronological and palaeoenvironmental frameworks within which to interpret the archaeological record has also inspired research in (initially) less promising regions of the world that have previously received little attention, such as the deserts of the Arabian Peninsula (e.g. Groucutt et al., 2015; Stimpson et al., 2015, 2016). The resulting palaeontological data have the potential to be interpreted on a variety of scales, ranging from considerations of changing climatic conditions during a single interglacial in a given region (e.g. Candy et al., 2016) to comparisons between different interglacials (e.g. Limondin-Lozouet and Preece, 2014) and even identification of patterns at the marine oxygen isotope substage level (Schreve, 2001b; Candy and Schreve, 2007; Ashton et al., 2008; White et al., 2013). Key to such studies have been important advances in the application of independent geochronological methodologies to Quaternary fluvial archives (reviewed in detail by Rixhon et al., 2017). Advances in radiometric dating techniques, such as optically stimulated luminescence (OSL), electron spin resonance (ESR) and uranium-series have provided crucial chronological control (e.g., Rittenour, 2008; Kock et al., 2009; Voinchet et al., 2015;). In tectonically-active areas, the presence of interbedded volcanic deposits can provide further opportunities for constraining the age of the fluvial sediments through the application of argon–argon dating (e.g. Maddy et al., 2012; Marra et al., 2016). In addition, amino-acid racemization dating (AAR) based on the calcitic opercula of the freshwater gastropod genus *Bithynia* has been shown to be more reliable than previous AAR methodologies, allowing the development of a

robust aminostratigraphy for the British Quaternary with potential for application in other regions (Penkman et al., 2007, 2011, 2013; cf. Westaway, 2009).

2. NW and Central Europe

A substantial body of research has been undertaken over the last decade in this important region, particularly in southern Britain and northern France. This is partly a feature of preservation potential; the unequal global distribution of fossiliferous fluvial sediments corresponds closely with that of calcareous bedrock outcrops, which promote fossil preservation, limiting detailed biostratigraphical research to regions dominated by limestone and chalk. Calcareous tufas have provided an important additional source of data (Dabkowski, 2014), albeit from sediment sequences usually outside major fluvial systems. Tufa sequences have been recorded across Europe (Pentecost, 1995; Sancho et al., 2015), but are particularly well represented in northern France and Britain, where they have mainly been attributed to MIS 11, MIS 5e and the Holocene (Dabowski, 2014; Limondin-Lozouet and Preece, 2014). The Triassic Muschelkalk outcrop in Germany also gives rise to notable occurrences of calcareous spring deposits formed on former subaerial floodplains, although these are generally more lithified than their Anglo-French counterparts and have thus generally been termed travertines. Important multiple travertines are interbedded with the terrace deposits of the Ilm at Weimar (Schreve and Bridgland, 2002) and the Wipper at Bilzingsleben (Mania, 1995). Although tufa and travertine deposits tend to be highly localized, they can preserve fossils that rarely survive in other fluvial sedimentary settings; this wide variety of palaeontological data is therefore particularly suitable for both biostratigraphy and palaeoclimatic reconstructions. As well as molluscs and vertebrates, plant remains (sometimes in the form of imprints of rapidly-encrusted leaves or fruits) are also common, although pollen is rarely well preserved (Dabkowski, 2014). The calcareous nature of tufas and travertines also means that they are suitable for a range of geochemical analyses (e.g. Dabkowski et al. 2012, 2015), and appropriate for radiometric dating using U-series techniques (e.g. Candy and Schreve, 2007; Sierralta et al., 2010).

2.1 Britain

Quaternary palaeontology in Britain has benefitted from government funding of multidisciplinary research projects through bodies such as English Heritage (now Historic England) and schemes such as the Aggregates Levy Sustainability Fund (ALSF). The English Rivers Palaeolithic Survey (TERPS), which commenced in 1991, resulted in an important baseline archive for Pleistocene archaeological research (Wymer, 1999), but provided only basic information pertaining to the palaeontological evidence that accompanied Palaeolithic assemblages. More detailed reviews of regional Pleistocene fossil records were subsequently provided by projects funded by the ALSF, such as the Trent Valley Palaeolithic Project (TVPP), which conducted an exhaustive review of both the palaeontological and archaeological records of the English Midlands (Schreve, 2007; Bridgland et al., 2014), and the Medway Valley Palaeolithic Project (MVPP), which conducted similar research in Kent (Briant et al., 2012; Chauhan et al, this volume). These projects were able to study fossiliferous localities that lacked archaeological material, adding valuable palaeontological and palaeoclimatic data to the underlying TERPS dataset. An overview of all the ALSF-funded projects was provided by White (2016). Additional data was accumulated during the lifetime of the Ancient Human Occupation of

Britain Project (AHOB), funded by the Leverhulme Trust, which undertook research at numerous fluvial localities (Ashton et al., 2011; see <http://ahobproject.org/database>).

The data accumulated by these research projects have continued to test and reinforce the biostratigraphical model developed by Schreve (2001a; Fig. 2), which proposed a succession of Mammal Assemblage-Zones (MAZs) representing post-Anglian (MIS 12) interglacials. An important aspect of recent research has been the recognition of greater climatic complexity recorded in terrestrial interglacial sequences, leading to a renewed appreciation of the potential ability of multi-proxy palaeontological analyses to discern marine oxygen isotope substages (e.g. Schreve, 2001b; Ashton et al., 2008; Koutsodendris et al., 2010, 2011; White et al., 2013; Candy et al., 2014, 2016). In Britain and northern Europe, this has been especially relevant to sequences attributed to the MIS 11 Hoxnian/Holsteinian Interglacial (e.g. Preece et al., 2007; Ashton et al., 2008; Candy et al., 2014), which is particularly well represented in the terrestrial record.

2.1.1 The Lower Thames

The terrace deposits of the River Thames, particularly in its lower reach, represent one of the most important terrestrial archives of the Middle and Late Pleistocene in the world. In addition to key MIS 9 and MIS 7 sites published earlier in the lifetime of FLAG (Schreve et al., 2007 and references therein), new data have been obtained from several important Lower Thames localities over the last decade, including several MIS 11 sites: Dierden's Pit, Swanscombe (White et al., 2013), Southfleet Road, Swanscombe (Wenban-Smith, 2013), East Hyde, Tillingham (Roe, 2001; White, 2012) and Clacton-on-Sea (White, 2012). Further work has also been undertaken on the MIS 9 sequence at Purfleet (Bridgland et al., 2013) and other MIS 9 localities downstream in eastern Essex (Roe and Preece, 2011; Roe et al., 2009, 2011), providing further insight into the differentiation of MIS 11 and MIS 9 in the British terrestrial record (Bridgland et al., 2001; Thomas, 2001; Roe et al., 2009). Over the wider Thames valley, mammalian fossils have assisted with the correlation of the Upper and Middle Thames terraces (Bridgland and Schreve, 2009).

Insert Figure 2 hereabouts

At Swanscombe, the biostratigraphical and palaeogeographical significance of the 'Rhenish' suite of freshwater molluscs, an important feature of the MIS 11 faunal succession in the Lower Thames, has been firmly established by new evidence from Dierden's Pit (White et al., 2013). The Swanscombe sequence, together with its downstream correlatives at East Hyde and Clacton, indicates that the six 'Rhenish' species did not colonize the Thames simultaneously, but appeared in a distinctive sequence that can be tied to particular stages in the development of the vegetation. At both Swanscombe and Clacton, 'Rhenish' taxa are largely absent from the earliest fluvial deposits, with the pioneer species *Pisidium clessini* and *Theodoxus danubialis* first appearing during pollen zone Ho II. These, together with *Belgrandia marginata*, become established at the onset of Ho III, just before the first evidence for estuarine conditions (in the form of brackish indicator species) appears at both sites. They are followed by *Corbicula fluminalis*, which appears during Ho IIIb, and *Borysthenia naticina*, which is absent from Clacton but appears in significant numbers slightly later than *Corbicula* at both Swanscombe and East Hyde (White et al., 2013). The timing of the first appearance of the final 'Rhenish' species, *Viviparus diluvianus*, is

less clear-cut, since it is a rare component of the Swanscombe fauna (Kerney, 1971) and is only represented by derived shells at Clacton. However, at East Hyde it appears to be present from early in the sequence, suggesting it was established in the lower Thames by at least pollen zone Ho IIIa (White et al., 2013). It has therefore been possible, on the basis of multiple strands of fossil evidence, to correlate MIS 11 fluvial sequences in the Lower Thames and to establish the timing of sea-level change and the migration of the estuarine environment during that interglacial (White et al., 2013; Fig. 3). This has shown that sea-levels rose relatively late in the interglacial, during Ho III, with a significant period during which the Thames was confluent with continental rivers such as the Scheldt; this was presumably also true of other fluvial systems flowing into the southern North Sea basin. This palaeogeographical situation allowed distinctive groups of invertebrate and vertebrate species (including humans) to colonize Britain (White et al., 2013).

Insert Figure 3 hereabouts

Other significant recent advances in knowledge of the Lower Thames sequence have arisen from work undertaken as part of developer-funded archaeological appraisal, particularly that related to construction of the high-speed railway connection to the Channel Tunnel (e.g. Bridgland et al., 2013; Wenban-Smith 2013) and within the remit of work funded by the ALSF. The construction of the Channel Tunnel Rail Link led to the unexpected discovery of an elephant-butchery site, preserved within deposits of a south-bank Thames tributary at Southfleet Road, Swanscombe (Wenban-Smith et al., 2006; Wenban-Smith, 2013). The carcass of an extinct straight-tusked elephant (*Palaeoloxodon antiquus*) was surrounded by an undisturbed scatter of flint tools (Wenban-Smith, 2015). Fossil assemblages recovered from the Southfleet Road sequence included vertebrates, molluscs, ostracods and pollen; the absence of 'Rhenish' molluscs indicates that the sequence pre-dates the Middle Gravels at Swanscombe and is therefore probably equivalent to the Lower Loam at Barnfield Pit, a conclusion consistent with the presence of Clactonian artefacts (Wenban-Smith, 2013). The vertebrate assemblage is strikingly similar to those from Swanscombe (Basal Gravel-Lower Loam) and Clacton, both of which have been assigned to the Swanscombe MAZ (Schreve, 2001a).

Work at the important MIS 9 sequence at Purfleet, undertaken as part of various developer-funded projects, allowed assessment of the palaeontological assemblages from the site, together with new analyses of stable isotopes and geochronology (Bridgland et al., 2013). This work confirmed the attribution of the interglacial to MIS 9 through the application of OSL and AAR and contributed new palaeoecological information. The MIS 9 interglacial has also been studied at various sites downstream in eastern Essex (Roe and Preece, 2011; Roe et al., 2011), allowing enhancement of knowledge of the evolution of the highstand MIS 9e estuary in the Lower Thames valley. Further south, the MVPP produced amino acid racemization data from several fossiliferous localities in eastern Essex, including the MIS 11 sequences at Clacton-on-Sea, East Hyde, Bradwell Hall, the MIS 9 sequences at Shoeburyness, Cudmore Grove, East Wick and the Last Interglacial (MIS 5e) site at East Mersea. The famous Levallois site at Baker's Hole, located in the Ebbsfleet Valley, NW Kent, has also been the subject of recent research following the construction of the Channel Tunnel Rail Link and Ebbsfleet International station (Scott et al., 2010; Wenban-Smith, 2014). This locality, dated to MIS 7, represents the

last surviving remnant of what was (before quarrying) a wide area containing Levalloisian lithic remains in undisturbed primary context, associated with assemblages of large mammals, molluscs, small vertebrates and ostracods; publication of detailed biostratigraphical and palaeoenvironmental analyses is in progress.

2.1.2 The River Trent and its tributaries

The Trent, Britain's third longest river, had until recently received considerably less attention than other British fluvial systems. Significant new insights into the origin and evolution of the Trent system have resulted from the Trent Valley Palaeolithic Project (TVPP), which conducted an exhaustive review of the geological, palaeontological and archaeological records of the English Midlands (Howard et al., 2007; White et al., 2007, 2010; Bridgland et al., 2014, 2015). The dating of the Trent terraces has been underpinned by biostratigraphical data, together with aminostratigraphic and OSL dating programmes and uplift modelling (Penkman et al., 2011, 2013; Bridgland et al., 2014; Westaway et al., 2015; Westaway, 2017). Pleistocene fossil assemblages from the Trent catchment are both less common and less well known than equivalent records from the aforementioned Lower Thames sites and adjacent catchments, such as the Warwickshire-Worcestershire Avon. The upper and middle reaches of the Trent include few calcareous rocks, fossils being preserved there only under exceptional circumstances, such as within the Allenton Terrace deposits south-west of Derby (Bridgland et al., 2014) and the floodplain gravels at Whitemoor Haye (Schreve et al., 2013). Both of these sites are located at the confluences of tributary rivers with the main Trent. At Allenton, calcareous gravels enriched by Carboniferous limestone clasts from upstream in the tributary Derwent valley are responsible for the localized preservation of vertebrate assemblages that include hippopotamus, on the basis of which these deposits have been attributed to MIS 5e (Bridgland et al., 2014). The younger Whitemoor Haye locality, at the Tame-Trent confluence, is situated within low-lying 'floodplain terrace' deposits, dated by radiocarbon and OSL to around 41–43 k cal a BP, placing them within the Middle Devensian (MIS 3; Schreve et al., 2013). This site is prone to waterlogging, another effective means of preserving organic remains; palaeotemperature reconstructions, based on beetle and chironomid assemblages, have indicated mean July temperatures of +8 to +11 °C and mean January temperatures of between –22 and –16 °C (Schreve et al., 2013).

The most significant fossil assemblages from the Trent catchment have been recovered from its lower reach, where numerous sites have now been recorded in the Balderton–Southrey Terrace of the Trent and in the Fulbeck and Tattershall terraces of its tributaries, the Witham and Bain, respectively (Bridgland et al., 2014, 2015; Fig. 4). A significant aspect of the late Middle and Late Pleistocene record in the Trent is the absence of deposits belonging to the MIS 11 and MIS 9 interglacials, which are well represented in fluvial systems further south, most notably the Thames. This has been attributed to extensive glaciation of the English Midlands during MIS 8 (White et al., 2010, 2017; Bridgland et al., 2014). The oldest fossiliferous sediments known from the Trent system are therefore those preserved within the Balderton-Southrey terrace and its equivalents in the Witham and Bain valleys (Brandon and Sumbler, 1988, 1991; Schreve, 2007; Bridgland et al., 2014). The Balderton Terrace of the Lower Trent is predominantly a cold-climate aggradation, from which characteristic fossils such as *Mammuthus primigenius* (woolly mammoth), *Coelodonta antiquitatis* (woolly rhinoceros) and *Ovibos moschatus* (musk ox) have been recovered (Brandon

and Sumbler, 1988, 1991; Bridgland et al., 2014). Earlier studies of these vertebrate assemblages also recognised the presence of interglacial species (Brandon and Sumbler, 1988, 1991; Lister and Brandon, 1991), which were subsequently established to have been derived from highly fossiliferous fine-grained deposits preserved low down in the sequence, typically filling channels cut into the Lias Clay bedrock or the lowermost part of the Balderton Formation (Bridgland et al., 2014). The fossil assemblages from these basal channels are representative of interglacial conditions and include molluscs, ostracods, small vertebrates, pollen, plant macrofossils and insects (White et al., 2007; Bridgland et al., 2014). The best record to date has come from Norton Bottoms, where significant exposures of fine-grained organic-rich sediments were recorded between 2006 and 2008 (White et al., 2007; White, 2012; Bridgland et al., 2014). Amongst the large mammal remains was a virtually complete cranium and horn cores of an adult aurochs (*Bos primigenius*), together with well-preserved molluscan assemblages (some specimens retaining their periostracum) that included the bivalves *Corbicula fluminalis* and *Pisidium clessini* and the land snail *Candidula crayfordensis*, none of which are known in Britain after MIS 7 (see Bridgland et al., 2014). Insect assemblages from Norton Bottoms are also a close match with those from British MIS 7 sites, lacking the exotic components that characterize MIS 9 and MIS 5e (Coope, 2001; Murton et al., 2001; Green et al., 2006; Bridgland et al., 2014); these have provided mutual climatic range (MCR) estimates for July temperatures (T^{max}) of +16 to +18°C and January temperatures (T^{min}) of -11 to +5°C (Bridgland et al., 2014). Complementary data derived from ostracod assemblages using the mutual ostracod temperature range (MOTR) method (Horne, 2007; Horne et al., 2012) have provided similar temperature estimates of T^{max} +15 to +21°C and T^{min} -4 and +3°C (Bridgland et al., 2014).

Insert Figure 4 hereabouts

Other sequences containing fine-grained sediments have been recorded in the Witham Valley, notably from boreholes at Coronation Farm and Stainfield (Bridgland et al., 2014). These yielded molluscs, pollen, plant macrofossil and insect remains; the molluscan assemblages contained no biostratigraphically-significant species, but the presence of *Bithynia* opercula allowed AAR dating to be applied, which suggested an age within MIS 7 (Penkman, 2007; Bridgland et al., 2014). In the valley of the River Bain, a left-bank tributary of the Trent-Witham system, quarrying at Tattershall Thorpe revealed interglacial sediments that contained fossils not found in deposits attributed to MIS 5e (Holyoak and Preece, 1985 and references therein), providing some of the earliest indications that both MIS 7 and MIS 5e were preserved in the terrace deposits in that area. The Last Interglacial (Ipswichian, MIS 5e) is represented in three parts of the Trent system, all of them areas where the main river is joined by a significant tributary. In the Middle Trent, sites at the confluence of the River Derwent with the Trent, at Boulton Moor and Allenton, have yielded hippopotamus fossils, together with beetles and plant remains (Arnold-Bemrose and Deeley, 1896; Jones and Stanley, 1974, 1975). In the Witham Valley, hippopotamus was also found at several sites in the vicinity of Fulbeck (Brandon and Sumbler, 1988, 1991; Howard et al., 1999; Bridgland et al., 2014), and further downstream MIS 5e sediments (without hippopotamus) were found at Tattershall Castle (Holyoak and Preece, 1985; Bridgland et al., 2014).

As well as providing age constraints for the terrace deposits, the new recognition of widespread MIS 7 interglacial deposits in the Lower Trent has been critical in the

identification of a late Middle Pleistocene (post-Anglian–pre-Devensian) glaciation in Britain during MIS 8, a considerable rarity globally. Indeed, widespread biostratigraphical evidence for MIS 7 deposits in Britain as far north as Bielsbeck in East Yorkshire appear to exclude the possibility of an eastern British glaciation south of the River Humber during MIS 6 (White et al., 2010, 2017). No interglacial sequences attributable to MIS 11 or 9 have been identified in the Trent system, providing evidence (albeit negative) for widespread destruction of this part of the late Middle Pleistocene record (White et al., 2010, 2017). Dating evidence from the Trent fluvial archive, in the form of biostratigraphy and aminostratigraphy, has been critical in constraining the age of this glaciation, demonstrating the utility of fluvial records where they interdigitate with glacial deposits (see Cordier et al., this issue).

2.1.3 The Fenland rivers

The rivers flowing into the basin of the Wash in eastern England were recently studied as part of another ALSF project (the Fenland Rivers of Cambridgeshire Project), which concluded in 2008 (Boreham et al., 2010). These included the Witham (before the latest Pleistocene this was the lower Trent; see Bridgland et al., 2014, 2015), Welland, Nene, Great Ouse and Cam, all of which have yielded well-preserved fossil assemblages. Morphostratigraphical approaches to fluvial deposits within the Fen Basin are extremely difficult to apply, since this is an area that has undergone very little uplift, resulting in poor differentiation of the Pleistocene terraces. Biostratigraphical approaches have also been attempted (Bridgland and Schreve, 2001), although the precise correlations of several key interglacial deposits in the Fen Basin with the marine oxygen isotope stage record remain controversial due to mixed or inadequate biostratigraphical signals (Boreham et al., 2010). This is largely due to the preservation of sediments representing more than one interglacial in close proximity beneath a single terrace surface (Boreham et al., 2010). Research in the Peterborough area has revealed considerable complexity in the fluvial deposits preserved there (Langford and Briant, 2004; White et al., 2010, 2016; Bridgland et al., 2014; Langford et al., 2014a, b). Once again, occurrences of *Corbicula fluminalis* and *Hippopotamus amphibius* have been key to distinguishing the Last Interglacial (Ipswichian, MIS 5e) from earlier interglacials in the Wash fluvial systems. The resulting synthesis of data enabled common patterns in these archives to be discerned, but also highlighted many significant differences between these rivers, all of which have shared a downstream valley during periods of low sea-level.

2.2 France

Multidisciplinary research projects investigating fluvial archives in NW France have largely focussed on the Somme and Seine river valleys. Over the last decade, several important palaeontological datasets from these areas have provided significant chronological and palaeoenvironmental information relevant to understanding Pleistocene climates and human occupation (see Chauhan et al., this issue). The French fluvial record is well-dated and includes several interglacial sequences particularly rich in diverse fossil groups that allow palaeoclimatic comparisons with nearby British records and other datasets at a European scale (e.g. Limondin-Lozouet and Preece, 2014). Alongside mammalian evidence (see Auguste, 2009), molluscan evidence remains a key element of French biostratigraphical schemes. The Lateglacial–Holocene molluscan successions recorded at numerous fluvial sites have been central to understanding the palaeoenvironmental significance of Pleistocene

vegetational changes and climatic conditions (Limondin-Lozouet, 2011). During transitional climatic phases, the stages of recolonization by molluscan faunas show common features, such as the occurrence of specific taxa and well-defined biotope successions. These malacological characteristics have been correlated with climatic phases and botanical evolution. For example, in the early Holocene of NW Europe the replacement of the landsnail *Discus ruderatus* by its congener *Discus rotundatus* has been linked to the spread of *Corylus* (Preece and Day, 1994; Limondin-Lozouet et al., 2005). The same faunal succession has also been identified within Pleistocene interglacial sequences at La Celle in the Seine Valley, dated to MIS 11, and at Caours in the Somme basin, dated to MIS 5e. Although pollen was not preserved at either of these sites, the first appearance of *D. rotundatus* can be used to identify the development of deciduous forest, replacing the pioneer open forest biotope (Limondin-Lozouet, 2011; Fig. 5).

Insert Figure 5 hereabouts

2.2.1 The Somme and Nord Pas-de-Calais

Research undertaken over the last 20 years on the fluvial terraces and loess sequences of the Somme Basin, and on the interactions between human populations and changing environmental conditions, has adopted interdisciplinary approaches for the analysis of Quaternary sequences and associated Palaeolithic sites (Antoine et al., 2003, 2007, 2010; Bahain et al., 2007, 2010; Bridgland et al., 2006). These studies, mainly targeting fluvial sequences, have highlighted the impact of cyclic climatic changes on sedimentation and river morphology, and especially the role of the 100 ka climatic cycles for the last million years, giving rise to stepped terrace formation (Antoine et al., 2007). Interglacial climatic optima are recorded by calcareous tufa sequences, which are especially well preserved in the cases of MIS 11 and 5e.

At the Carrière Carpentier site at Abbeville, sediments (the 'White Marl') dated to MIS 15 using ESR (584 ± 48 ka) and their stratigraphical position within the Somme terrace system have yielded molluscs, large vertebrates and small mammals (Antoine et al., 2015, 2016; Voinchet et al., 2015). A noteworthy occurrence within the small vertebrate faunas are rodent molars tentatively attributed to *Arvicola cantiana/mosbachensis*. The presence of this taxon is indicative of a younger age in the early Middle Pleistocene, i.e. younger than MIS19-17 (Antoine et al., 2016). The large mammal fauna includes the Hundsheim rhinoceros (*Stephanorhinus hundsheimensis*), wild boar (*Sus scrofa priscus*) and red deer (*Cervus elaphus*), representing the Cromerian faunal association known from West Runton and Pakefield. When compared with British Cromer Forest-bed sites, the mammalian fauna from Carrière Carpentier appears to be contemporaneous with Pakefield, more recent than West Runton, but older than the Boxgrove fauna (Antoine et al., 2016). The Carrière Carpentier sequence is within Alluvial Formation VII of the Somme system (Antoine, 1994, 2000) and is associated with two other sites located in the same terrace, Carrière Léon and Moulin Quignon (Bahain et al., 2016). Characteristic freshwater mollusc species from this alluvial formation include *Tanousia* cf. *stenostoma*, *Borysthenia naticina* and *Bithynia troschelii*, which form an assemblage typical of Cromerian interglacial fluvial deposits in NW Europe (Meijer and Preece, 1996; Preece, 2001). Specimens of *T. cf. stenostoma* from Moulin Quignon are similar to those of Little Oakley (Bahain et al., 2016). This occurrence represents the first record of this species in France; in Britain it is unknown after MIS 13.

Recent archaeological rescue excavations in the Somme valley have led to the discovery of several alluvial sequences containing molluscan faunas (Locht et al., 2013). At Rue Boileau in Amiens, a sequence exposed in a pit adjacent to the famous site of Saint Acheul includes a white silt horizon that has been correlated on morphostratigraphical grounds with the Saint Acheul tufa; the latter has been dated to MIS 11 on the basis of a characteristic '*Lyrodiscus* assemblage' of land snails, supported by U/Th dating (Limondin-Lozouet and Antoine, 2006). The molluscan fauna from Rue Boileau contains several critical species typical of this assemblage, together with several xerophilous taxa (*Cernuella virgata*, *Candidula unifasciata*) previously unknown from St Acheul (Locht et al., 2013). This assemblage has therefore been tentatively attributed to a later phase of the interglacial, post-dating the climatic optimum, when heavily-forested environments were being replaced by more open habitats characterised by xerophilous land snails (Limondin-Lozouet et al., 2015).

The important fluvial sequence preserved at Caours represents the only Eemian (MIS 5e) interglacial sequence in the Somme basin (Antoine et al., 2006). Overlying a periglacial gravel bed allocated to MIS 6 (Alluvial Formation I of the Somme system), the fluvial fine-grained sequence consists of calcareous silts and tufa deposits including thin organic layers, from which a wealth of palaeontological material has been recovered, most notably non-marine molluscs. These have provided a record of the glacial-interglacial transition between MIS 6 and MIS 5. Cold-tolerant faunas similar to those of the Younger Dryas, including some boreo-alpine taxa (*Columella columella*), were replaced by numerous thermophilous taxa, highlighting a hiatus corresponding to the very earliest part of the interglacial (Limondin-Lozouet, 2011). Following this, the malacological succession indicates the development of forest environments during an early part of MIS 5, followed by the climatic optimum of MIS 5e and a subsequent cooling (Limondin-Lozouet and Preece, 2014). Several species of Central European and Mediterranean origin (e.g. *Daudebardia rufa*, *Ruthenica filograna*, *Platyla polita*) occurred during the Eemian climatic optimum and constitute the first record of a regional biostratigraphical marker for the last interglacial (Limondin-Lozouet and Preece, 2014; Fig. 5). In addition, the Caours site has yielded archaeological material, providing an important record of human occupation in northern France during the Last Interglacial. It has now been identified as a butchery site, with lithic industries directly associated with mammalian remains (Auguste, 2009). A further important aspect of the Caours sequence is evidence for palaeomagnetic reversal thought to represent the Blake Event; identification of this geomagnetic excursion, which has been correlated with the start of the continental Eemian Stage, indicates that the palaeoclimatic and archaeological records from Caours post-date the MIS 5e interglacial peak (Sier et al., 2015).

To the north of the Somme basin, in the neighbouring region of Nord Pas-de-Calais, a fluvial sequence was discovered in 2012 at Waziers (Hérisson et al., 2015). This sequence comprises several peat layers overlying fluvial silts and sands, deposited in a large meandering system. Preliminary geomorphological and palaeontological observations, including the presence of loess covering the fluvial deposits and the peat and the occurrence of assemblages of interglacial mammals (aurochs and red deer) and aquatic molluscs (*Belgrandia marginata*, *Anisus septemgyratus*) suggested an Eemian age for this sequence. This was corroborated by a minimum age of 103 ± 3.5/-3.4 ka obtained by U/Th dating of calcareous charophyte oogonia, extracted

from a fine-grained tufa layer directly underlying the peat (Hérisson et al., 2015). This site is only the second to provide evidence of Neanderthal occupation during the Eemian in Northern France.

Finally, the late MIS 5 fluvial sequence at Ailly-sur-Noye, located ~30 km south of Amiens, includes three Palaeolithic levels (Locht et al., 2013). The oldest lithic industries and associated mammals (*Bos primigenius*, *Equus* sp.) lie at the top of a basal fluvial gravel attributed to the late Saalian/Eemian on the basis of its geomorphological position within the terrace system. The upper part of the sequence, consists of coarse gravels (also containing Palaeolithic artefacts), covered by a thin calcareous tufa crust with facies typical of interglacial tufa deposits like those described at Caours and likely to represent MIS 5e. Above this, the sediments consist of fine grey silts capped by a calcareous mud containing two further archaeological levels characterized by Levallois material. Molluscan faunas from these deposits are 'Arianta' assemblages typical of transitional climatic phases (Puisségur, 1976); further evidence for deteriorating climatic conditions is provided by a decline in the numbers of the aquatic snail *Belgrandia marginata*, a well-known thermophilous species typical of Pleistocene interglacial sequences that occurs in abundance at Caours (Antoine et al., 2006). The molluscan evidence indicates a transitional context from an interglacial phase to an early glacial episode, thought to represent the period spanning MIS 5d to MIS 5a (Locht et al., 2013); this is the first regional malacological record for this time period in France.

2.2.2 The Seine

New research undertaken at the site of Saint-Pierre-lès-Elbeuf has provided additional evidence for the age and palaeontological content of the important sequence there (Cliquet et al., 2009). Above the lowermost palaeosol (Elbeuf IV) lies a white sand and a calcareous tufa from which a characteristic 'Lyrodiscus assemblage' was obtained: this fauna is a regional biostratigraphical marker correlated with the MIS 11 interglacial. Systematic malacological sampling within this unit has demonstrated a high level of homogeneity within the molluscan population, which is largely dominated by forest land snails (up to 80 % of the total shells). This implies that the tufa accumulated relatively rapidly during the optimum phase of MIS 11. New IRSL dates obtained from the Elbeuf IV soil (475 ± 38 ka) and on the white sand (396 ± 32 ka) confirm the MIS 11 correlation (Cliquet et al., 2009).

The Upper Seine valley preserves another significant tufa sequence at the site of La Celle, which has been known for more than a century as an important source of non-marine molluscs and plant macrofossil impressions. New research at this site has been undertaken in order to improve understanding of the palaeontological assemblages and to provide additional chronological control, which has been correlated with MIS 11 (Limondin-Lozouet et al., 2006, 2010). Dating evidence has been provided by the geomorphological position of the site, within an old terrace of the Seine deposited prior to the 'Nappe de Soucy' of the Yonne valley system dated at 350–300 ka (Limondin-Lozouet et al., 2006), together with the occurrence of land snails characteristic of the 'Lyrodiscus fauna', known from several French and British tufa sequences of MIS 11 age (Rousseau et al., 1992; Limondin-Lozouet and Antoine, 2006; Preece et al., 2007). Additional age control has now been provided by radiometric dating (ESR/U-series on tooth enamel of horse and ESR quartz), which

have produced a mean age of ~400 ka (Bahain et al., 2010; Voinchet et al., 2015). However, although this geochronological evidence indicates an age equivalent to MIS 11, the existing methods do not allow precise timing within the interglacial. The La Celle tufa provides the longest known MIS 11 malacological succession in NW Europe, which has been used to generate detailed reconstructions of the development of forest cover (Limondin-Lozouet et al., 2010, 2015) and as a biostratigraphical standard against which other MIS 11 malacological successions can be compared (Limondin-Lozouet et al., 2015). Successive appearances of forest species, first from an Atlantic corridor and later from central and southern Europe, allow characterization of vegetational development. The initial immigrants indicate the spread of closed habitats, while a peak in the diversity of thermophilous snails corresponds to the optimum phase of deciduous forest expansion. A subsequent decrease in forest species, together with an expansion of hygrophilous taxa, indicates the decline of closed canopy conditions. The occurrence of critical forest species, especially those now extinct or occurring far beyond their modern ranges, provides a framework within which molluscan successions from other tufa sites in northern France and southern Britain can be understood (Fig. 6).

Insert Fig. 6 hereabouts

Leaf impressions from La Celle have allowed the identification of 24 taxa, including some Mediterranean plants such as *Buxus*, *Ficus* and *Celtis* (Limondin-Lozouet et al., 2010). Mammalian remains from La Celle (Auguste, 2009; Limondin-Lozouet et al., 2010) include *Macaca sylvanus* (Barbary macaque) and *Hippopotamus amphibius*; the former last occurred in Britain during MIS 9, whereas the latter was absent from Britain during MIS 11 (Schreve, 2001a); the occurrence of hippopotamus in northern France during MIS 11 is therefore of particular interest, perhaps indicating a biogeographical barrier that prevented this species colonizing Britain at that time.

2.3 Germany and Belgium

One of the key sites in this region is the multiperiod locality at Schöningen, Germany, the evidence from which was summarized by Schreve et al. (2007). Debate on the age of this sequence has continued over the last decade and was recently summarized by Urban and Bigga (2015). As is the case in Britain, the occurrence of *Theodoxus danubialis* has been attributed biostratigraphical significance in Germany, where *T. serratilineiformis* (= *danubialis*) has been suggested to indicate an MIS 11 age (e.g. Meng and Wansa, 2005, 2008). This is by no means certain; in the middle Neckar valley (SW Germany), an exposure of fluvial gravels at the site of Bietigheim-Bissingen, ~20 m above the modern River Enz, yielded a molluscan fauna including *Theodoxus serratilineiformis* and *Cochlostoma scalarinum saueri* (Bibus and Rähle, 2003). On the basis of terrace stratigraphy and molluscan biostratigraphy, these deposits were dated to MIS 13, although this age is not accepted by Meng (2007).

In the southern North Sea Basin the western coastal plain of Belgium preserves a complex sequence of Pleistocene marine and fluvio-estuarine deposits. Recent analysis of brackish and freshwater fossil assemblages (including ostracods, pollen and Foraminifera) recovered from boreholes in this region have indicated deposition near the upper tidal limit of an estuary (Bogemans et al., 2016). Of biostratigraphical and biogeographical significance is the occurrence, in the Zoutenaai core, of a single valve of the freshwater ostracod *Scottia browniana*, which is unknown in the nearby

British record after MIS 11 (Whittaker and Horne, 2009). Given that the oldest of the Belgian channel fills are suggested to date to MIS 9 (Bogemans et al., 2016; cf. Vanhoorne, 2003), it is possible that this fossil has been reworked; however, its presence might also indicate that *S. browniana* became extinct in continental Europe later than in Britain. This might well be expected, since the fluctuating island status of Britain from MIS 12 onwards enhances turnover in faunal composition and biogeographical differences.

3. Eastern Europe and Russia

Faunal evidence in this region has been recovered from the significant fluvial archives preserved in the valleys of the south-flowing rivers of the East European Platform, the Dniester, Dnieper, Don and Volga, and has been key to unravelling their age and evolution (Matoshko et al., 2002, 2004; Schreve et al., 2007; Bridgland and Westaway, 2008, 2014). To the south-west of the Russian Plain, several fossiliferous localities are known within the Prut and Danube basins, although these are predominantly represented by lacustrine sequences (see Markova and van Kolfschoten, 2012). Several important Middle Pleistocene faunal localities are known from the Dnieper, mostly recovered from fluvial deposits assigned to Terrace IV, correlated with the Likhvin Interglacial (=MIS 11) on the basis of pollen and mammalian biostratigraphy (Markova, 2006; Markova and van Kolfschoten, 2012; Fig. 7). Amongst the best studied of these are the mammal assemblages from Gunki, Pivikha and Chigirin (Fig. 7). The Gunki site has also yielded a molluscan fauna, considered to represent the Early Euxinian (Markova and van Kolfschoten, 2012). In the Don and Desna basins, the earliest small mammal faunas have been correlated with the early Middle Pleistocene (Markova, 2007). Here, faunal assemblages have been correlated with both interglacial and glacial stages (Fig. 7). The biostratigraphical significance of several mollusc species from the Don Basin was highlighted by Kondrashov (2007), who noted that the occurrence of species such as *Borysthenia intermedia*, *Lithoglyphus jahni* and *Viviparus fasciatus* allows correlation of the Don basin faunas with the early Middle Pleistocene Cromerian faunas of Western Europe. In the Volga basin, a small mammal fauna that includes *Arvicola cantiana*, *Lagurus transiens-lagurus* and *Clethrionomys rufocanus* has been described from a fluvial sequence at Rybnaya Sloboda, at the mouth of the right-bank tributary Kama River, and has been attributed to the Likhvin Interglacial on biostratigraphical grounds (Markova and van Kolfschoten, 2012; Fig. 7). Similar faunas have been reported from sites in the middle and lower Volga, at Chernyi Yar and Spasskoe (Markova and van Kolfschoten, 2012). To these can be added a considerable amount of vertebrate, molluscan and vegetational data that has been collected since the 1960s in the southern Urals region, summarized in a flurry of papers published since 2007 detailing the records of the Belaya and Lemeza rivers (e.g. Chlachula, 2010; Danukalova et al., 2007, 2008, 2009, 2011, 2016; Puchkov and Danukalova, 2009; Yanina, 2013; Yakovlev et al., 2013). Changes to the fauna and flora in this area appear to have been more muted than the adjacent northwestern territories due to the absence of local glaciers during some cold stages (Danukalova et al., 2009).

Insert Figure 7 hereabouts

Further afield, in Ukraine, mollusc and ostracod assemblages have been described from a sequence of Early–early Middle Pleistocene fluvial gravels and loams exposed in a quarry at Skala Podil'ska, in the valley of the River Dniester (Boguckij et al.,

2009). The freshwater mollusc assemblages from the basal fluvial gravels (Unit I) include *Theodoxus serratilineiformis* (= *danubialis*) and *Viviparus lungershauseni* (= *fasciatus*; Kondrashov, 2007). Also present was the ostracod *Limnocythere tuberculata*, which is also indicative of the Early Pleistocene (Dykan, 2003). The overlying loams of Unit III also contained *T. danubialis*, together with *Lithoglyphus neumayri*. The ostracod fauna from this upper unit contained species characteristic of colder water bodies (Boguckij et al., 2009). Palaeoclimatic research in Russia has also been significantly advanced by the development of the QUINSIB database, which contains details of over 600 fossil insect localities (Kuzmina, 2014). The development of this resource has allowed large amounts of unpublished data and published sources in Russian to be made widely accessible.

4. Southern Europe and Iberia

4.1 Spain and Portugal

A surge of interest in Iberian fluvial systems followed the successful FLAG Biennial Meeting at Castelo Branco in 2010, particularly with the development of improved chronological frameworks for several fluvial systems in Spain and Portugal (such as the Tagus/Tejo, Minho, Douro/Duero, Mondego and Guadiana rivers), which have been considerably refined through the application of new luminescence techniques (Cunha et al., 2008; Vis et al., 2008; Martins et al., 2010; Antón et al., 2012; Ramos et al., 2012; Viveen et al., 2012, 2013; Carvalhido et al., 2014; Sancho et al., 2016). However, with the exception of a few notable regions previously summarized by Schreve et al. (2007), preservation of palaeontological material is generally poor across the Iberian Peninsula and the terraces of these rivers have yielded relatively little new fossil material (e.g. Cunha et al., 2012). Fluvial archives such as that preserved in the Ter River basin, northeastern Spain, have proved difficult to date due to the absence of fossiliferous deposits (Garcia, 2015). A notable recent discovery is the site of Barranc de la Boella (Catalonia, Spain), where vertebrate remains (including a butchered elephant) were recovered in association with stone tools from deposits of the Francolí river (Vallverdú et al., 2014). Dating of this site was based on palaeomagnetic and cosmogenic determinations, supported by biostratigraphical evidence derived from the vertebrate faunas, which included *Mimomys savini* and *Mammuthus meridionalis*, suggesting a late Early Pleistocene age (Mosquera et al., 2015).

In central Spain, faunal assemblages have been recovered from the sites of Pinedo and Cien Fanegas near Toledo, in the +25–30 m terrace of the River Tajo (Tagus), which have also yielded Acheulian archaeology. These gravels been dated to between 290 and 220 ka using AAR and luminescence (pIR–IRSL) techniques, spanning a period between MIS 9 and 7 (López-Recio et al., 2015). The gravel pit at Pinedo has yielded fossils of straight-tusked elephant, together with hippopotamus, rhinoceros, deer, horse and bovids (López-Recio et al., 2015). Elsewhere in the Middle Tajo, large mammal assemblages have been recovered in association with stone tools from the 40 m terrace at Toledo; these include *Mammuthus trogontherii*, *Equus caballus*, *Hippopotamus amphibius*, *Megaloceros savini*, *Eliomys quercinus*, *Allocricetus bursae*, *Microtus brecciensis* and *Apodemus sylvaticus*, all characteristic of the Middle Pleistocene (Sesé et al., 2000). Palaeoclimatic studies from this region include reconstructions based on herpetofaunal assemblages from three Spanish

localities dated to MIS 11, including the fluvial deposits at Áridos-1, in the valley of the River Jarama SE of Madrid, and the fluvio-lacustrine sequence at Ambrona (Blain et al., 2015). These herpetofaunal assemblages include numerous taxa of high environmental and climatic sensitivity, which can be used as valuable indicators of palaeoclimate, through the application of MCR analyses (e.g. Martínez-Solano and Sanchiz, 2005; Blain et al., 2008).

4.2 Italy

The Pleistocene vertebrate succession in the Italian peninsula is relatively well known (e.g. Gliozzi et al., 1997), although much of this evidence has been recovered from cave and fissure-fill sequences in karstic regions (e.g. Sardella et al., 2003; Pandolfi and Petronio, 2011a, b; Pandolfi et al., 2013). The potential relationships between these isolated cave sequences and local fluvial systems have not yet been explored in detail, but research in other regions suggests that biostratigraphical schemes can be usefully developed in this way (e.g. Yang et al., 2011; Bridgland et al., 2014, Westaway, 2016). In the mountainous regions of northern and central Italy, fossil assemblages have also been obtained from lacustrine sequences, with fluvio-deltaic deposits sometimes represented (e.g. Girotti et al., 2003; Limondin-Lozouet et al., 2017). Arguably the best known Italian fluvial archives are those from the Tiber River basin, the record for which spans much of the Pleistocene. The palaeo-Tiber and its tributaries have been extensively studied in the area around Rome (Caloi et al., 1998; Di Stefano et al., 1998; Milli et al., 2004; Petronio et al., 2011). Recent work on this system has been undertaken by Marra et al. (2014), who identified six biochronological units (Slivia, Ponte Galeria, Isernia, Fontana Ranuccio, Torre in Pietra and Vitinia), spanning a period from ~600 ka to at least MIS 7, although this scheme has been criticised (Sardella et al., 2015; cf. Marra et al., 2015).

In central Italy, fossiliferous fluvial sequences representing a period spanning the Early Pleistocene to the early Middle Pleistocene have been recorded in trenches and boreholes at several localities in the Anagni Basin (Bellucci et al., 2012, 2014). The important Coste San Giacomo locality, known since the late 1970s, has yielded a Villafranchian large mammal fauna, including taxa such as *Mammuthus meridionalis*, *Hippopotamus* sp., *Equus stononis*, *Gazella borbonica*, *Sus strozzi* and *Homotherium* sp. More recently, microfaunal assemblages obtained from the Coste San Giacomo 1 core, drilled in 2009, have allowed more detailed palaeoecological reconstructions to be attempted (Bellucci et al., 2012, 2014). Analyses of small vertebrate, ostracod, pollen and charophyte assemblages, together with sedimentological data, indicate an initially forested landscape giving way to an open alluvial plain fed by sand-bed rivers (Bellucci et al., 2014). In addition, the ostracod assemblage contained sufficient extant taxa to attempt summer and winter air temperature reconstructions using the MOTR method, providing estimates of mean July temperatures of between 15°C and 22°C and mean January temperatures of between -5°C and 5.2°C. However, it should be noted that the assemblage used for these estimates was derived from 8 separate assemblages recovered from a large section of the borehole (Bellucci et al., 2014) and may therefore represent an averaging of a long period of time. Small vertebrate assemblages included the biostratigraphically significant Early Pleistocene vole *Mimomys pliocaenicus*; this species, together with evidence from pollen biostratigraphy and magnetostratigraphy, suggests a Gelasian age for the Coste San

Giacomo faunal unit, falling somewhere between 2.2 and 1.95 Ma (Bellucci et al., 2014).

5. The Levant and the Arabian Peninsula

Fossil preservation in the Levant is relatively poor, despite the widespread presence of calcareous bedrock. It has been suggested that the formation of calcreted gravels, a characteristic of such semi-arid regions (e.g. Candy et al., 2004, 2005), might be responsible for the loss of faunal remains - in the Upper Orontes, for example, cemented Pleistocene terrace deposits appear to have undergone repeated decalcification and re-cementation, resulting in the weathering-out of calcareous clasts and, presumably, calcareous fossils (Bridgland et al., 2012). The general absence of biostratigraphical data from fluvial sequences (although see Tchernov, 1981, 1994 for reviews of data from cave and lake sequences in Israel) is mitigated by the presence of Pleistocene lava flows interbedded within the terrace sequences of many of the Turkish and Syrian fluvial systems that can be dated using various radiometric techniques (Sharkov et al., 1998; Bridgland et al., 2007; Demir et al., 2007, 2012, Seyrek et al., 2008; Westaway et al., 2009; Maddy et al. 2012).

Beyond the Levant, in the interior of the now hyper-arid Arabian Peninsula, significant fluvial archives are largely absent; however, fossils have been recovered from lacustrine sequences in the Nefud and Rub' al Khali deserts (e.g. Thomas et al., 1998; Groucutt et al., 2014; Stimpson et al., 2014, 2016). Most of these assemblages have been dated to the Late Pleistocene (MIS 5e and younger) and Holocene, although in exceptional circumstances older assemblages have been preserved (see below, Stimpson et al., 2014, 2016). As new sites are discovered and recorded, there is increasing potential to develop biostratigraphical schemes linking Arabian lake sites to Levantine fluvial systems.

5.1 The Levant

Productive new research in Syria, investigating the terraces of the River Orontes (Bridgland et al., 2003, 2012) and on the catchments of Mesopotamian rivers such as the Euphrates and Tigris (Demir et al., 2007, 2008, 2012; Westaway et al., 2009) has been curtailed in recent years due to ongoing conflicts in this war-torn region. The summary of research provided by Schreve et al. (2007) therefore remains largely up to date, with the exception of a handful of recently-published papers detailing research undertaken before 2009 (e.g. Bridgland et al., 2012). Fossiliferous sequences are also relatively rare in accessible reaches of these rivers in Turkey (e.g. Demir et al., 2007, 2008, 2012; Westaway et al., 2009; Seyrek et al., 2014a, b). Vertebrate faunas have provided important biostratigraphical evidence constraining the ages of the Orontes terraces (Bridgland et al., 2003, 2012; Bridgland and Westaway, 2007; Mishra et al., 2007). Biostratigraphical evidence provided by the vertebrate assemblages from Latamneh has necessitated the reattribution of the Orontes QfIII terrace in the Middle Orontes to an age of 1.2–0.9 Ma, leading to a revision of the age model for the Orontes terrace sequence (Bridgland et al., 2012; cf. 2003). Given this revision, it is no longer tenable to attribute the Middle Orontes terraces to formation in response to 100 ka Milankovitch climatic forcing (cf. Bridgland and Westaway, 2008b). A revised model for the evolution of that reach of the Orontes, resulting from this modification of the supposed age of the key biostratigraphical marker, sees similarities between the incision history in the Hama –

Latamneh area and those determined from the Euphrates in its southern Turkish and Syrian reaches, related to the crustal characteristics of the Arabian Platform (Bridgland et al., 2017). Downstream from Latamneh, molluscan and ostracod faunas were obtained from river-cliff locations at Karkour, alongside the Orontes channel as it traverses the subsiding Ghab Basin, including the ostracod *Cyprideis torosa*. This taxon is generally associated with brackish environments, developing noded valves in salinities below ~5‰, but can also tolerate hypersaline conditions in lakes and water bodies prone to desiccation, which is presumably how it comes to be in the Ghab sediments. The faunas here also include freshwater elements, both ostracods and molluscs, the latter including the large viviparid gastropod *Apameaus apameae* (Bridgland et al., 2012), which is also recorded at the key comparator (biostratigraphically somewhat younger) Israeli locality at Gesher Benot Ya'aqov, in the Jordan Valley, where it is an index fossil for the definition of the 'upper freshwater series' or 'Viviparus Beds' of the Benot Ya'aqov Formation (Picard, 1963; Tchernov, 1973; Goren-Inbar and Belitzky, 1989; Bar-Yosef and Belmaker, 2010). The last appearance of *Apameus apamae* in the Jordan Valley was at ~ 240 ka, on the basis of U-series dating (Kafri et al., 1983; Moshkovitz and Magaritz, 1987; Heller, 2007); its presence at two localities in the Ghab basin, Syria (Bridgland et al., 2012) and at the site of Alaattin Köyü in Turkey (Seyrek et al., 2014) therefore suggests a latest Early Pleistocene to late Middle Pleistocene age for these deposits.

5.2 The Arabian Peninsula

In recent years, the Arabian Peninsula (Saudi Arabia, Yemen, Oman and the United Arab Emirates) has emerged as an important region for research into low-latitude Quaternary environmental change (e.g. Petraglia, 2007; Parker, 2009; Armitage et al., 2011; Groucutt and Petraglia, 2012; Delagnes et al., 2012). Expansions of plant and animal communities, including humans, into and through the Arabian interior have occurred on several occasions, corresponding with humid climatic phases during the Pleistocene, when savanna-type landscapes prevailed in what are now hyper-arid regions (Vaks et al., 2007, 2013; Rosenberg et al., 2011, 2013; Breeze et al., 2015; Jennings et al., 2015a, 2017; Parton et al., 2015a). However, the highly fragmentary nature of Arabian terrestrial sequences, compounded by poor stratigraphical and chronological control on associated palaeontological assemblages, has hampered the construction of biostratigraphical and palaeoenvironmental frameworks.

A major hindrance to better regional understanding of the Arabian record is the lack of perennial fluvial systems and their resulting sedimentary archives; evidence for fluvial activity consists largely of poorly-dated gravels and alluvial deposits resulting from ephemeral and highly seasonal catchments (for recent reviews see Breeze et al., 2015, 2016). The principal drainage across central Arabia consists of several eastward-flowing wadis that presently carry water only seasonally. It is likely that these systems were repeatedly activated during humid periods in the Pleistocene and early Holocene (Powers et al., 1966; Chapman, 1971; Anton, 1984; Edgell, 2006), although some are now choked in places by dunes (Holm, 1960), suggesting that they have been inactive as continuous systems for a considerable period of time. At a smaller scale, more localized alluvial fans have formed around the bases of steep-sided jebels and dykes, from which intermittent streams flowed during wet phases (e.g. Parton et al., 2015b; Jennings et al., 2015b). It has been suggested that most of the trans-Arabian wadi systems were incised in their current configurations by the early Quaternary, based on relationships between basal gravel deposits and dated lava

flows (Al-Sayari and Zötl, 1978; Anton, 1984). Younger terraces have also been dated using early radiocarbon techniques (e.g. Jado and Zötl, 1984) but, given the unreliability of these methods, these ages require verification. More recent dating programmes have applied OSL and U-series techniques to fluvial deposits in south-central Arabia (e.g. Maizels, 1987, 1990; Blechschmidt et al., 2009; McLaren et al., 2009; Parton et al., 2010, 2013; Rose et al., 2011; Sitzia et al., 2012; Atkinson et al., 2013), contributing to an increasingly detailed chronology for the Late Pleistocene and Early Holocene. However, the general absence of substantial fluvial archives in Arabia is reinforced by the fact that the most significant stratified archaeological assemblages reported in recent years have been found in association with either lacustrine sequences, representing a source of fresh water, or raw materials suitable for stone tool production (e.g. Armitage et al., 2011; Delagnes et al., 2013; Hilbert et al., 2014; Groucutt et al., 2015; Jennings et al., 2015b; Scerri et al., 2015).

6. Data from other regions

Beyond the 'core regions' of NW and Central Europe, palaeontological datasets from fluvial sequences are increasingly used as a basis for regional biostratigraphies and palaeoenvironmental reconstructions. At the time of the last review (Schreve et al., 2007), these consisted entirely of mammalian data; since then, significant new information, including invertebrate and floral assemblages, have been published and are included in the summaries below.

6.1 China and the Far East

In China, the Nihewan Formation, comprising fluvio-lacustrine sediments containing abundant mammalian fossils, is widely distributed in the Nihewan Basin of northern Hebei Province. This region has yielded the densest concentration of Early Pleistocene Palaeolithic sites outside Africa (Dennell, 2013), together with fossil assemblages (vertebrates, molluscs, pollen) that have long been considered to be of Early Pleistocene age, although alternative ages ranging from late Pliocene to Middle or Late Pleistocene have also been proposed (Zhao et al., 2010). ESR dating of quartz obtained from the Majuangou and Banshan sites provided ages of between 1.70 and 1.35 Ma (Liu et al., 2014). Further south in the Three Gorges region, located in the transitional zone between the upper and middle reaches of the Yangtze (Changjiang) River, several fossiliferous fluvial sequences have now been reported (Pei et al., 2013). Seven fluvial terraces (T₇–T₁) have been identified and dated using ESR, TL, OSL and radiocarbon techniques (Pei et al., 2013). The site at Jingshuiwan, assigned to T₂ and dated to 75.9 ± 3.7 to 64.5 ± 4.1 ka using OSL, yielded an assemblage of 58 mammalian fossils including *Stegodon orientalis* (Pei et al., 2010).

In northwest Hunan Province, the terraces of the Suoxi River have been dated using ESR and TL techniques, allowing them to be correlated with neighbouring karstic cave systems (Yang et al., 2011). Although none of the dated localities in this study proved to be fossiliferous, presumably due to the local sandstone bedrock, such research highlights the potential for fluvial archives to provide chronological evidence that can be linked to calcareous sequences with potential to provide palaeontological data. Palaeoclimatic research in East Asia using methods such as beetle MCR estimates has been hampered by a lack of knowledge of modern distributions of beetle species and the sparse distribution of meteorological stations, especially in Siberia (Shiyake, 2014). Japan is exceptional in having conditions that have allowed

application of the Mutual Climatic Range (MCR) method, having a dense meteorological observation network and higher quality distribution data for its extant beetle fauna (Shiyake, 2014). Experimental work at the site of Nojiriko, Nagano Prefecture has provided the first MCR reconstructions from East Asia (Shiyake, 2014).

6.2 South Asia

The Indian subcontinent has yielded a range of palaeoecological and biostratigraphical data, in the form of vertebrate and invertebrate fossil assemblages, pollen records and archaeological material, much of which has been recovered from fluvial sediments of the Narmada, Godavari, Manjra, Son, Ghod, Krishna and Mahanadi rivers (Chauhan, 2008). Most of these deposits have been assigned ages no older than the Middle Pleistocene, although some formations in the Narmada Valley might date to the Early Pleistocene (Tiwari and Bhai, 1997; Tiwari, 2001). The application of modern dating techniques has shown that many of these sedimentary units have the potential to be significantly older or younger than previously thought (Chauhan 2008 and references therein), a situation that is also prevalent in the Levant (Bridgland et al., 2012) and Arabia (e.g. Thomas et al., 1998, contra Stimpson et al., 2014, 2015). Until reliable chronological frameworks are available for these fluvial archives, their usefulness in terms of regional biostratigraphy remains somewhat limited.

The Siwalik deposits of northern India, Pakistan, Nepal and Myanmar represent one of the best-studied fluvial sequences in Asia. The Upper Siwaliks have been divided into three subdivisions, the youngest two of which (the Pinjor and Boulder Conglomerate Formations) represent most of the Quaternary, spanning the period from 2.58 to 0.2 Ma (Prasad, 2001; Mishra et al., 2010). Mammalian fossils are abundant within the Pinjor Formation (2.58 to 0.6 Ma), which has yielded at least 98 vertebrate species including *Elephas hysudricus*, *Stegodon insignis*, *Rhinoceros*, *Sivatherium*, *Equus sivalensis*, *Bos acutifrons* and *Cervus palaeindicus*, together with carnivores such as *Canis pinjorensis*, *Crocota felina* and *Pachycrocota brevirostris* (Nanda, 2002, 2008). However, the stratigraphic range of most of these species within the Pinjor Formation is unknown, limiting their biostratigraphical application (Dennell et al., 2008). Nevertheless, reconstructions of the palaeoecology of the fluvial landscapes within which the Pinjor Formation was deposited have been attempted (Dennell et al., 2008), including predator-prey interactions and the circumstances under which fossil assemblages were accumulated and buried.

Two younger faunal horizons, roughly equivalent to the late Middle and Upper Pleistocene respectively, are preserved on the Indo-Gangetic plain in India and in Sri Lanka (Nanda, 2008; Mishra et al., 2010), although these have yielded only 16 and 26 mammalian taxa, respectively. These 'post-Siwalik' faunas suggest that a large proportion of the Upper Siwalik Pinjor fauna became extinct during the Middle Pleistocene. Because the stratigraphical relationships between these numerous fluvial localities remain unclear, it has not yet been possible to propose a testable biostratigraphical framework for the region. However, a relative chronostratigraphic, biostratigraphic and archaeological correlation for the Narmada Basin formations has been proposed (Badam, 2007; Patnaik et al., 2009). At Hathnora, assemblages of vertebrates (including herpetiles and fish), molluscs, ostracods, charophytes and

pollen, in association with archaeological material, have been reported (Patnaik, 2000; Patnaik et al., 2009). Freshwater molluscs have also been recovered from the sites at Bhedaghat and Devakachar (Hirdepur Formation), including 18 species of gastropods and bivalves (Kotlia and Joshi, 2011). Several lines of evidence suggest that Southeast Asia was dominated by a mosaic of savannah, open woodland and evergreen forest throughout much of the Pleistocene, leading to suggestions that this region served as a refugium for hominins and other mammal species during glacial periods (Louys and Turner, 2012)

Well-dated fossiliferous sequences in Southeast Asia are relatively rare and are predominantly preserved in caves, although the syntheses described here include some data from fluvial deposits. Broad characteristics of the extinctions of Pleistocene large vertebrates in this region were recently reported by Louys et al. (2007), together with consideration of the palaeoenvironmental requirements of many of these species (Louys and Meijaard, 2010). Many species which became extinct appear to have been endemic to specific areas, with others suffering severe range reduction before their eventual demise. Members of the latter group include proboscideans (*Stegodon* and *Palaeloxodon*), the pygmy hippopotamus (*Hexaprotodon*), the orangutan (*Pongo*), hyenas (*Crocuta* and *Hyaena*), the giant panda (*Ailuropoda*), tapirs (*Tapirus* and *Megatapirus*), rhinoceroses (*Rhinoceros*), and the giant Asian ape, *Gigantopithecus*. The loss of these species is likely to have been the result of a combination of climatic changes (Louys et al., 2007) and human impacts (Corlett, 2007). Unlike other regions which experienced megafauna extinctions, such as South America (see below), eustatic changes in sea level in Southeast Asia seems to have been an important factor (Louys et al., 2007).

On the island of Java, Indonesia, fluvial deposits of the Solo River have recently yielded both vertebrate fossils and archaeological material. An extensive survey of the region around Matar, close to the site at Ngandong (where fossils of *Homo erectus* were recovered in the early 1930s), revealed the presence of at least three river terraces; palaeontological and archaeological assemblages were recovered from the First and Second terraces (Fauzi et al., 2016). Tentative comparisons with other Pleistocene faunal localities on Java suggest that the Matar assemblage is younger than the Middle Pleistocene Kedungbrubus Fauna but older than the Punung Fauna, since it contains several species (such as *Stegodon trigonocephalus*, *Bubalus paleokarabau*, *Bibos paleosondaicus* and *Hexaprotodon sivalensis*) that occur in the former but not in the latter (Fauzi et al., 2016). The Punung Fauna, the type locality for which is Punung Cave, Indonesia, has been dated to 128 ± 15 and 118 ± 3 ka using luminescence and U-series techniques (Westaway et al., 2007). Further work is required in order to understand fully the regional biostratigraphy.

Insert Fig. 8 hereabouts

6.3 North and South America

A significant proportion of Pleistocene palaeontological research on these continents has been driven by a desire to understand the reasons for megafaunal extinctions (e.g. Faith et al., 2009; Haynes, 2009; Scott, 2010; Doughty et al., 2013) and the timing of the first human colonization of the Americas (e.g. Goebel et al., 2008; Rothammer and Dillehay, 2009; Pitblado, 2011), as well as dynamics between the two (e.g. Gill et al., 2009). Fluvial archives are also potential sources of palaeontological evidence

relevant to debates surrounding the Great American Biotic Interchange (GABI), a period of faunal exchanges between North and South America. The isolation and faunal endemism of South America ended with the formation of the Panamanian land bridge and the onset of the GABI, with the first of four major faunal exchanges now thought to coincide with the base of the Pleistocene at ~2.6 Ma (Reguero et al., 2007; Woodburne, 2010). Further major pulses, at 1.8, 0.7 and 0.125 Ma (Fig. 8) appear to have coincided with periods of lower sea-level, leading to the suggestion that expanded coastal regions, cooler climates and associated changes to local flora promoted animal dispersals at these times (Woodburne, 2010).

Insert Fig. 9 hereabouts

There are few Pleistocene sites in South America from which well-dated fossil material has been recovered from secure stratigraphical contexts; correlations between localities have therefore frequently been based on the biostratigraphical scheme proposed for the fossiliferous beds of the Pampean region of Argentina, which has been periodically updated over the last two decades (e.g. Cione and Tonni, 1999, 2001, 2005). The Quaternary mammal faunas of South America have also been reviewed from time to time by several authors (see Prado and Alberdi, 2009 and references therein). Major phases of mammalian dispersal occurred during the Pleistocene Ensenadan and Lujanian biochrons (Fig. 9), which have also been dated on the basis of the Pampean sequence (Cione and Tonni, 1999, 2001, 2005). Biostratigraphical significance has been ascribed to four species of *Neosclerocalyptus* (Glyptodontidae) which occurred during the Pleistocene in the Pampean region of Argentina (Zurita et al., 2009a). Two of these, *Neosclerocalyptus pseudornatus* and *N. ornatus* are indicative of the Ensenadan Stage, the former occurring between 1.07 and 0.98 Ma and the latter between 0.98 and 0.40 Ma (Fig. 9). Two other species, *N. gouldi* and *N. paskoensis*, are thought to represent the Bonaerian and Lujanian stages, respectively (Zurita et al., 2009a; Fig. 9). This biostratigraphical scheme has therefore become a standard with which other South America sequences are now routinely compared (e.g. Lopes et al., 2010; Tonni et al., 2009; Beilinson et al., 2015).

In Bolivia, the Tarija Valley preserves some of the richest Pleistocene mammal localities in South America, although the dating of these assemblages remains unclear (Coltorti et al., 2007; Tonni et al., 2009; Zurita et al., 2009b). A Middle Pleistocene age has been suggested on the basis of mammalian biostratigraphy (e.g. Tonni et al., 2009 and references therein), although they have also been attributed to a much younger period on the basis of radiocarbon dating (~44 – 21 ka BP). A similar lack of reliable dating for fossiliferous sites has hampered the bio- and chronostratigraphical correlation of South American Pleistocene faunas elsewhere; in Brazil, Lopes *et al.* (2010) published ESR dates of between 226 and 34 ka for a vertebrate assemblage from Chuí Creek; this broad age range, spanning a time period encompassing the final part of MIS 7 until MIS 4, is probably due to the reworked nature of the Chuí Creek vertebrate assemblages (Pereira et al., 2012). In Venezuela, strata such as the Taima Taima fossil bed and fluvial sites at Muaco, Cucuruchú, and Quebrada Ocando have been dated on the basis of biostratigraphy (Carlini et al., 2008).

In Uruguay, the fossiliferous beds of the Sopas Formation have yielded vertebrate faunas, together with freshwater molluscs, plant remains and other trace fossils (Ubilla et al., 2004, 2009, 2016; Ubilla and Martínez, 2016). The mammalian

assemblage includes extinct taxa, such as the capybara *Neochoerus aesopi*, the glyptodont *Neuryurus rudis*, and two extinct species of deer (*Antifer ultra* and *Morenelaphus brachyceros*). This assemblage has been correlated on the basis of biostratigraphy with the Lujanian Stage (Late Pleistocene–Early Holocene) of the Pampean region of Argentina; this is supported by radiocarbon AMS dates from the vertebrate material range from $39,900 \pm 1,100$ to $33,560 \pm 700$ BP (cal 45,389 to 42,025 ya) and TL/OSL ages derived from the sediments range from $71,400 \pm 11,000$ to $27,400 \pm 3,300$, indicating a MIS 3 age for the fauna (Ubilla et al., 2016). Palaeoenvironmental evidence suggests that the Sopas formation represents open habitats, savannahs and woodlands (Ubilla et al., 2016).

Insert Fig. 10 hereabouts

6.4 Australia

In Australia, Quaternary palaeoenvironmental research based on fluvial archives has a long pedigree in two important regions of the continent: the arid interior of the Lake Eyre Basin, where fluvial, lacustrine and aeolian sequences representing the last ~300 ka are preserved (recently reviewed in detail by Habeck-Fardy and Nanson, 2014), and the extensive meandering river systems of southeastern Australia, particularly the well-dated terraces of the Lachlan and Macquarie rivers in the Murray-Darling Basin (Kemp and Spooner, 2007; Yonge and Hesse, 2009; Kemp and Rhodes, 2010). Faunal and floral responses to Pleistocene climate change in these regions remain less well understood, primarily due to the rarity of stratified fossil assemblages and significant issues with directly dating fossil material (Price et al., 2013; Westaway et al., 2017). As is the case in North America, much research in Australia has been focussed on the extinction of megafaunal species during the Late Pleistocene; approximately 96 % of the large mammal fauna was extinct by ~45 ka, a period broadly concurrent with human colonisation, although the extent to which these extinctions can be directly related to human activity remains a source of considerable debate (e.g. Koch and Barnosky, 2006; Prideaux et al., 2007, 2010; Price et al., 2015; Dortch et al., 2016; Johnson, 2016; Johnson et al., 2016; Westaway et al., 2017; Fig. 11). Overhunting and the burning of the landscape by people, with a corresponding abrupt reduction in plant diversity, have been suggested as causal factors (Porch and Kershaw, 2010; Rule et al., 2012), but significant climate change in the period between 50 and 46 ka, resulting in a major shift to more arid conditions and a corresponding drop in water levels in Lake Eyre and Lake Frome, is also considered to have played an important role (Murphy et al., 2011; Cohen et al., 2012, 2015; Sakaguchi et al., 2013).

In the Lake Eyre Basin, fossils of 21 megafaunal mammal species have now been recorded (Webb, 2008, 2009), including a giant wombat-like marsupial (*Diprotodon*), giant short-faced kangaroo (*Procoptodon goliah*), a large flightless bird (*Genyornis newtoni*), giant goanna (*Varanus priscus*) and the 'marsupial lion' (*Thylacoleo carnifex*). In addition, smaller mammal species have been recovered from the upper Katipiri Formation (MIS 6–4), including the Southern brown bandicoot (*Isodon obesulus*), the extinct Eastern hare wallaby (*Lagorchestes leporides*), the Pale field rat (*Rattus tunneyi*) and the Western grey kangaroo (*Macropus fuliginosus*); the fossil occurrences of the three extant species are well beyond their modern distributions (Webb, 2009). However, the lack of chronological control for much of this fossil material means that the first and last appearances of many mammal species in the

wider Lake Eyre Basin remain poorly understood. Better biostratigraphical evidence has been forthcoming in eastern Australia, where vertebrate and molluscan assemblages (44 taxa in total) have been recovered from the Darling Downs, within fluvial deposits of the Kings Creek (Price and Sobbe, 2005; Price et al., 2011). Palaeoenvironmental analyses have indicated that a mosaic of local habitats, including vine thickets, scrublands and open grasslands, prevailed during the late Pleistocene; increasing aridity led to a contraction of the more wooded environments in favour of grassland. Dating of the Kings Creek sequences using a variety of techniques (OSL, U-series and radiocarbon) has shown that individual megafaunal species responded independently to climatic and environmental change, revealing a more complex staggered extinction pattern in southeastern Australia prior to the arrival of humans (Price et al., 2011; Fig. 11).

Insert Fig. 11 Australian biostrat chart

7. Synthesis

Regional biostratigraphies based on mammalian and molluscan assemblages continue to contribute significantly to the dating and correlation of Pleistocene fluvial deposits on a global scale. However, the quality of these datasets varies widely, depending on the potential for preservation of fossils in a given region and the priority given to palaeontological and biostratigraphical research. The period since the 2007 review of regional biostratigraphies has seen a steady accumulation of data in many parts of the world. Those based on mammalian and molluscan assemblages, in particular, have continued to be tested and developed, especially in NW Europe, and clearly have a significant role to play in the dating and correlation of Quaternary deposits. Biostratigraphical frameworks are still predominantly based on mammals and other vertebrates, which is not surprising considering the robust nature of their fossils. Molluscs are also commonly used. Other groups, such as reptiles, amphibians, ostracods and insects are still only infrequently used as dating tools (e.g. Gleed-Owen, 1988, 1999, Coope, 2001; Griffiths, 2001; Whittaker and Horne, 2009; Borodin et al., 2013), but can provide invaluable complementary information as palaeoclimatic indicators. The relative rarity of non-mammalian vertebrate remains, which has been noted in fluvial sequences in regions such as Europe (Holman, 1998; Roe et al., 2009), Australia (Price and Sobbe, 2005) and Africa (Stoetzel et al., 2012) is usually attributed to taphonomic biases caused by the relative fragility of these fossils, although only a few studies specifically addressing this issue have been published (e.g. Pinto Llona and Andrews, 1996, 1998).

The predominance of work in the cooler temperate regions is in large part a result of the common preservation there of vertebrate and molluscan faunas in fluvial contexts that coincide with calcareous settings. In the warmer temperate Mediterranean region limestone abounds but dryland soil processes have led to reprecipitation of calcareous cements that seems to have coincided with the destruction of fossils. It is probably no accident that the key late Early Pleistocene fauna site of Latamneh in the Orontes in Syria is in one of the few reaches of that river with terrace gravels that are not calcareously cemented (Bridgland et al., 2012). The only other Orontes sites with significant faunas are preserved in the Ghab, where the presence of fossils might owe much to waterlogging in this fluvio-lacustrine subsiding basin (see Section 5).

There is an increasing need for research to address the validity of inferred patterns of faunal (including hominin) exchange between neighbouring regions. These have the potential to address important questions about the timing and nature of palaeoenvironmental change in response to climate change, by identifying key drivers of dispersal and the waxing and waning of biogeographical barriers. Of critical importance when comparing adjacent regions is the accuracy of chronological frameworks, which are required to establish whether significant palaeoenvironmental changes and faunal turnovers occurred synchronously in different regions, or whether leads and lags can be identified. It is often not possible to detect diachroneity in faunal change due to the lack of precision in radiometric methods, especially in the Early and Middle Pleistocene. Improved dating is thus required to determine patterns of north–south exchange between refugia (in the Northern Hemisphere), an issue of further relevance in respect of Latamneh, which has been suggested as a possible cold-stage assemblage that includes taxa represented in interglacials further north (Bridgland et al., 2012).

Detailed analyses are also important when it comes to reconstructing palaeoenvironmental change through longer fluvial sequences, with the potential to detect smaller-scale climatic oscillations, potentially correlateable with oxygen isotopic substages (e.g. Schreve, 2001b; White et al., 2013). These problems are well known to Quaternary biostratigraphers and palaeoecologists. The value of biostratigraphical and palaeoecological data has led to such studies becoming an increasingly common element of multi-disciplinary research; such projects are often driven primarily by archaeological research questions. Evidence corroborating and strengthening established biostratigraphical frameworks (Schreve, 2001a, b) has been obtained, together with substantial new understanding of the palaeoclimatic and palaeoecological significance of numerous plant and animal species that occur commonly within fossil assemblages (e.g. Candy et al., 2012, 2015, 2016).

Certain taxa appear to have been extremely widespread at particular times during the Pleistocene, only for their ranges to fragment in response to subsequent environmental pressures. The distribution of the hippopotamus (*Hippopotamus amphibius*) has long had significance in Britain, where it is a distinctive marker-fossil for the Last Interglacial (MIS 5e) (Sutcliffe, 1964; Currant and Jacobi, 2001; Schreve, 2001a, 2009). Similarly, distinctive freshwater mollusc species such as those of the 'Rhenish suite' characteristic of the MIS 11 Thames were clearly able to disperse rapidly across large parts of Europe. These two examples illustrate the potential for fluvial systems to connect otherwise separate biogeographic provinces, linking regional biostratigraphic schemes at certain times. Terrestrial species, for which large rivers are potential barriers to dispersal, are less likely to be able to profit from fluvial connectivity. However, in the case of the '*Lyrodiscus*' fauna that characterises molluscan faunas in Britain and northern France during MIS 11, some continuity in woodland habitat was clearly required to allow this biome to develop.

Some of the most significant patterns in the palaeontological record are evident at the extremities of regions and in differences between continental and island records. For example, in NW Europe the well established British biostratigraphical record differs from that of its nearest continental neighbours; both regions benefit from excellent preservation of fossils and well-dated fluvial stratigraphies, allowing direct

comparison of the faunas and floras of Pleistocene interglacials. The occurrence of hippopotamus in Britain and on the near-continent continues to be a point of difference, for example. This taxon occurs in abundance in Britain during the Late Pleistocene only during the Last Interglacial (MIS 5e, Ipswichian), whereas in northern France it has been found in assemblages dated to MIS 11 and MIS 7.

Palaeoclimatic reconstructions based on multiple fossil proxies have proved particularly useful for understanding interglacials, especially on the northern fringes of Europe. This sort of research has only been possible in regions where long, detailed sedimentary sequences are preserved. In Britain, much of this evidence has been derived from fossils of a diverse range of thermophilous flora and fauna that now occur in more southerly parts of continental Europe, or even further afield. At a basic level, qualitative measures of enhanced warmth (or cold) can be shown by the relative abundances of thermophilous or cryophilous species within fluvial deposits. More precise palaeotemperature estimates can be obtained from quantitative methods based on palaeoecological proxies, which now have a long pedigree within Quaternary research in northwest Europe (e.g. Atkinson et al., 1987; Zagwijn, 1996; Horne et al., 2012). Such approaches employ the known climatic tolerances, or the mapped climatic range, of extant plant or animal species in order to infer the warmest (Tmax) and coldest (Tmin) temperatures under which a given assemblage of species could survive (Candy et al., 2010). Fossils of these indicator species (or assemblages of groups of species) within Pleistocene fluvial deposits can thus be used as proxies for the palaeotemperature regime that prevailed at the time the deposit accumulated; within longer sequences, it is also possible to infer significant climatic changes from the fossil record.

Consideration of the ecological preferences of mollusc and ostracod species is important and can bolster arguments from faunal comparison data that can otherwise seem circular or overly reliant of negative (absence) evidence. An interesting observation from recent years has been the occurrence of halophytic taxa in warm-climate fluvio-lacustrine settings, where evaporation enhances salinity. This was noted in the reach of the Orontes in northern Syria, where the aforementioned subsiding Ghab basin is crossed (Bridgland et al., 2012). At that land-locked locality there is a low potential for confusion with an estuarine environment, but in more coastal settings the co-occurrence of certain species could raise problems for environmental interpretation.

8. Conclusions

Since the last major review of biostratigraphical data derived from Quaternary fluvial archives in 2007, significant new research on Pleistocene fossil assemblages has been undertaken on almost every continent. Established regional biostratigraphical frameworks have been challenged, modified and, in many cases, strengthened by the new data generated by this work. Over the 20-year lifespan of FLAG, much of the evidence summarized here has been obtained from fluvial sequences in the NW European 'hotspot' that includes Britain and France, together with important archives from Germany, Iberia and Italy and Eastern Europe. Research in Europe has included the reporting of new sites, dating or re-dating of known localities to improve chronological control, and the development of high-resolution palaeoenvironmental reconstructions based on fossil assemblages and geochemical analyses. There have also been important advances in understanding of the palaeoclimatic and

1324 palaeoecological significance of numerous plant and animal species that occur
1325 commonly within Pleistocene fossil assemblages; in combination with improved
1326 chronological frameworks, these provide important evidence for changing climate and
1327 environments through time.

1328
1329 In the Levant and South Asia, where the archaeological record has driven the research
1330 agenda, the recognition of the utility of fossil assemblages as chronological and
1331 palaeoenvironmental tools has led to them being increasingly studied, despite
1332 considerable taphonomic difficulties. The Arabian Peninsula, in particular, has been
1333 the subject of an increased focus of studies relating to hominin dispersals out of
1334 Africa, but because well preserved fluvial deposits are largely absent in this arid
1335 region, fossils have mainly been derived from lacustrine and cave sequences. The
1336 potential for developing biostratigraphical correlation schemes, linking the Arabian
1337 record to better-preserved fluvial archives in neighbouring regions such as the Levant,
1338 is therefore an exciting possibility for future research.

1339
1340 Europe and the Middle East have in common long-timescale records of human
1341 occupation; beyond these areas, in the Americas and Australia, human impacts came
1342 much later and have been studied mainly in the context of human contributions to the
1343 extinctions of Pleistocene megafaunas. The chronology of the Great American Biotic
1344 Interchange, together with refinements in South American biostratigraphical schemes
1345 in countries such as Brazil, Argentina and Bolivia continue to be major areas of
1346 research. Similarly, in Australia, frameworks of faunal extinctions have been
1347 developed. Australia has an apparently impoverished Middle and Late Pleistocene
1348 biostratigraphical record compared with other parts of the world, and many of the
1349 species that went extinct during this period are poorly dated. Archaeological sites
1350 attesting to potential human impact are rare and detailed ecological information for
1351 most extinct megafauna is lacking. As a result, the processes leading to megafaunal
1352 extinction remain unclear, although the weight of evidence points to a direct human
1353 impact as a major cause of extinction.

1354
1355 Although it is now possible to generate reliable radiometric dates for river terrace
1356 sequences in many parts of the world, enabling their correlation with the globally
1357 applicable marine oxygen isotope record, in some regions this remains extremely
1358 difficult. Where such robust chronological frameworks exist, providing fossils are
1359 also well preserved, it is possible to explore more detailed patterns in the occurrences
1360 of plant and animal species during the Pleistocene.

1361 **Acknowledgements**

1362 The authors wish to thank two anonymous reviewers for their constructive comments,
1363 which improved this manuscript.
1364

References:

- Al-Sayari, S.S., Zötl, J., 1978. *The Quaternary Period in Saudi Arabia, Vol.I: Sedimentological, Hydrogeological, Hydrochemical, Geomorphological, and Climatological Investigations in Central and Eastern Saudi Arabia*. Springer-Verlag, New York
- Antoine P. 1994. The Somme valley terrace system (northern France); a model of river response to Quaternary climatic variations since 800,000 BP. *Terra Nova* **6**, 453–464.
- Antoine, P., Rousseau, D.-D., Lautridou, J.-P., Hatté, C., 1999. The Last Interglacial-Glacial climatic cycle in loess-palaeosols successions of northwestern France. *Boreas* **4**, 551–563.
- Antoine, P., Lautridou, J.P., Laurent, M., 2000. Long-term fluvial archives in NW France: response of the Seine and Somme rivers to tectonic movements, climatic variations and sea level changes. *Geomorphology* **33**, 183–207.
- Antoine, P., Auguste, P., Bahain, J.-J., Coudret, P., Depaepe, P., Fagnart, J.-P., Falguères, C., Fontugne, M., Frechen, M., Hatté, C., Lamotte, A., Laurent, M., Limondin-Lozouet, N., Loch, J.-L., Mercier, N., Moigne, A.-M., Munaut, A.-V., Ponel, P., Rousseau, D.-D., 2003. Paléoenvironnements Pléistocènes et peuplements Paléolithiques dans le bassin de la Somme (nord de la France). *Bulletin de la Société Préhistorique de France* **100**, 5–29.
- Antoine, P., Limondin-Lozouet, N., Auguste, P., Loch, J.-L., Ghaleb, B., Reyss, J.-L., Escudé, E., Carbonel, P., Mercier, N., Bahain, J.-J., Falguères, C., Voinchet, P., 2006. Le tuf de Caours (Somme, France) : mise en évidence d’une séquence eemienne et d’un site paléolithique associé. *Quaternaire* **17**, 281–320.
- Antoine, P., Limondin-Lozouet, N., Chaussé, C., Lautridou, J.P., Pastre, J.F., Auguste, P., Bahain, J.J., Falguères, C., Ghaleb, B., 2007. Pleistocene fluvial terraces from northern France (Seine, Yonne, Somme): synthesis and new results. *Quaternary Science Reviews* **26**, 2701–2723.
- Antoine, P., Auguste, P., Bahain, J.-J., Chaussé, C., Falguères, C., Ghaleb, B., Limondin-Lozouet, N., Loch, J.-L., Voinchet, P., 2010. Chronostratigraphy and palaeoenvironments of Acheulean occupations in Northern France (Somme, Seine and Yonne valleys). *Quaternary International* **223-224**, 456–461.
- Antoine, P., Moncel, M.H., Loch, J.-L., Limondin-Lozouet, N., Auguste, P., Stoezel, E., Dabkowski, J., Voinchet, P., Bahain, J.-J. and Falgueres, C., 2015. Dating the earliest human occupation of Western Europe: new evidence from the fluvial terrace system of the Somme basin (Northern France). *Quaternary International*, **370**, 77–99.
- Antoine P., Moncel M.H., Limondin-Lozouet N., Loch J.L., Bahain J.J., Moreno D., Voinchet P., Auguste P., Stoezel E., Dabkowski J., Bello S.M., Parfitt S.A., Tombret O., Hardy B., 2016. Palaeoenvironment and dating of the Early Acheulean localities from the Somme River basin (Northern France): New discoveries from the High Terrace at Abbeville-Carrière Carpentier. *Quaternary Science Reviews*, **149**, 338–371
- Anton, D., 1984. Aspects of geomorphological evolution; palaeosols and dunes in Saudi Arabia. In: Jado, A., Zötl, J. (Eds.), *The Quaternary Period in Saudi*

- 1410 *Arabia, Vol.2: Sedimentological, Hydrogeological, Hydrochemical,*
 1411 *Geomorphological, and Climatological Investigations in Western Saudi Arabia.*
 1412 Springer-Verlag, New York, 275–296.
- 1413 Armitage, S.J., Jasim, S., Marks, A., Parker, A.G., Usik, V.I., Uerpmann, H., 2011.
 1414 The Southern route “Out of Africa”: evidence for an early expansion of modern
 1415 humans into Arabia. *Science* **331**, 453–456.
- 1416 Arnold-Bemrose, H.H., Deeley, R.M., 1896. Discovery of mammalian remains in the
 1417 old river gravels of the Derwent near Derby. *Quarterly Journal of the Geological*
 1418 *Society of London* **52**, 497–510.
- 1419 Ashton, N., Lewis, S.G., Parfitt, S.A., Penkman, K.E.H., Coope, G.R., 2008. New
 1420 evidence for complex climate change in MIS 11 from Hoxne, Suffolk, UK.
 1421 *Quaternary Science Reviews* **27**, 652–668.
- 1422 Ashton, N.M., Lewis, S.G., Stringer, C.B., 2011. *The Ancient Human Occupation of*
 1423 *Britain*. Developments in Quaternary Science **14**, Elsevier, Amsterdam.
- 1424 Atkinson, T.C., Briffa, K.R., Coope, G.R., 1987. Seasonal temperatures in Britain
 1425 during the last 22,000 years, reconstructed using beetle remains. *Nature* **325**,
 1426 587–592.
- 1427 Atkinson, O.A.C., Thomas, D.S.G., Parker, A.G., Goudie, A.S., 2013. Late
 1428 Quaternary humidity and aridity dynamics in the northeast Rub’ al-Khali, United
 1429 Arab Emirates: implications for early human dispersal and occupation of eastern
 1430 Arabia. *Quaternary International* **300**, 292–301.
- 1431 Auguste, P., 2009. Evolution des peuplements mammaliens en Europe du Nord-Ouest
 1432 durant le Pléistocène moyen et supérieur. Le cas de la France septentrionale.
 1433 *Quaternaire* **20**, 527–550.
- 1434 Badam, G.L., 2007. On the biostratigraphy and biochronology of the central Narmada
 1435 Valley Quaternary. In: Sankhyan, A.R., Rao, V.R. (Eds). *Human Origins,*
 1436 *Genome and People of India: Genomic, Palaeontological and Archaeological*
 1437 *Perspectives*. Anthropological Survey of India, Kolkata, 153–164.
- 1438 Bahain, J.-J., Falguères, C., Laurent, M., Voinchet, P., Dolo, J.-M., Antoine, P.,
 1439 Tuffreau, A., 2007. ESR chronology of the Somme River Terrace system and
 1440 first human settlements in Northern France. *Quaternary Geochronology* **2**, 356–
 1441 362.
- 1442 Bahain, J.-J., Falguères, C., Dolo, J.-M., Antoine, P., Auguste, P., Limondin-Lozouet,
 1443 N., Locht, J.-L., Tuffreau, A., Tissoux, H., Farkh, S., 2010. ESR/U-series dating
 1444 of teeth recovered from well-stratigraphically age-controlled sequences from
 1445 Northern France. *Quaternary Geochronology* **5**, 371–375.
- 1446 Bahain, J.-J., Limondin-Lozouet, N., Antoine, P., Voinchet, P., 2016. Réexamen du
 1447 contexte géologique, chrono et bio-stratigraphique du site de Moulin Quignon à
 1448 Abbeville (Vallée de la Somme, France). *L'Anthropologie* **120**, 344–368.
- 1449 Bates, M.R., Keen, D.H., Lautridou, J-P., 2003. Pleistocene marine and periglacial
 1450 deposits of the English Channel. *Journal of Quaternary Science* **18**, 319–337.
- 1451 Beaumont, P., 1999. *Northern Cape: INQUA XV International Conference Field*
 1452 *Guide*. McGregor Museum, Kimberley, South Africa.

- 1453 Beilinson, E., Gasparini, G.M., Soibelzon, L.H., Soibelzon, E., 2015. Insights into
1454 Pleistocene palaeoenvironments and biostratigraphy in southern Buenos Aires
1455 province (Argentina) from continental deposits. *Journal of South American Earth*
1456 *Sciences* **60**, 82–91.
- 1457 Belluci, L., Mazzini, I., Scardia, G., Bruni, L., Parenti, F., Segre, G.A., Naldini, E.S.,
1458 Sardella, R., 2012. The site of Coste San Giacomo (Early Pleistocene, central
1459 Italy): palaeoenvironmental analysis and biochronological overview. *Quaternary*
1460 *International* **267**, 30–39.
- 1461 Bellucci, L., Bona, F., Corrado, P., Magri, D., Mazzini, I., Parenti, F., Scardia, G.,
1462 Sardella, R., 2014. Evidence of late Gelasian dispersal of African fauna at Coste
1463 San Giacomo (Anagni Basin, central Italy): Early Pleistocene environments and
1464 the background of early human occupation in Europe. *Quaternary Science*
1465 *Reviews* **96**, 72–85.
- 1466 Bennike, O., Lindgård, E., Granat, H.J., Preece, R.C. and Viehberg, F., 2011. A new
1467 Middle Pleistocene interglacial sequence from Måløv, Sjælland, Denmark.
1468 *Geological Survey of Denmark and Greenland Bulletin* **23**, 29–32.
- 1469 Bibus, E., Rähle, W., 2003. Stratigraphische Untersuchungen an molluskenführenden
1470 Terrassensedimenten und ihren Deckschichten im mittleren Neckarbecken
1471 (Württemberg). *Eiszeitalter und Gegenwart* **53**, 94–113.
- 1472 Blain, H.A., Bailon, S., Cuenca-Bescós, G., 2008. Early–Middle Pleistocene
1473 palaeoenvironmental change based on the squamate reptile and amphibian
1474 proxies at the Gran Dolina site, Atapuerca, Spain. *Palaeogeography,*
1475 *Palaeoclimatology, Palaeoecology* **261**, 177–192.
- 1476 Blain, H.-A., Lozano-Fernandez, I., Olle, A., Rodríguez, J., Santonja, M., Pérez-
1477 González, A., 2015. The continental record of Marine Isotope Stage 11 (Middle
1478 Pleistocene) on the Iberian Peninsula characterized by herpetofaunal
1479 assemblages. *Journal of Quaternary Science* **30**, 667–678.
- 1480 Blechschmidt, I., Matter, A., Preusser, F., Riekezapp, D., 2009. Monsoon triggered
1481 formation of Quaternary alluvial megafans in the interior of Oman.
1482 *Geomorphology* **110**, 128–139.
- 1483 Bogemans, F., Roe, H.M., Baeteman, C., 2016. Incised Pleistocene valleys in the
1484 Western Belgium coastal plain: age, origins and implications for the evolution of
1485 the southern North Sea Basin. *Palaeogeography, Palaeoclimatology,*
1486 *Palaeoecology* **456**, 46–59.
- 1487 Boguckij, A.B., Łanczont, M., Łacka, B., Madeyska, T., Nawrocki, J., 2009.
1488 Quaternary sediment sequence at Skala Podil'ska, Dniester River basin (Ukraine):
1489 Preliminary results of multi-proxy analyses. *Quaternary International* **198**, 173–
1490 194.
- 1491 Boreham, S., White, T.S., Bridgland, D.R., Howard, A.J., White, M.J., 2010. The
1492 Quaternary history of the Wash fluvial network, UK. *Proceedings of the*
1493 *Geologists' Association* **121**, 393–409.
- 1494 Borodin, A., Markova, E., Zinovyev, E., Strukova, T., Fominykh, M., Zykov, S.,
1495 2013. Quaternary rodent and insect faunas of the Urals and Western Siberia:
1496 connection between Europe and Asia. *Quaternary International* **284** 132–150.

- 1497 Brandon, A., Sumblar, M.G., 1988. An Ipswichian fluvial deposit at Fulbeck,
1498 Lincolnshire and the chronology of the Trent terraces. *Journal of Quaternary*
1499 *Science* **3**, 127–133.
- 1500 Brandon, A., Sumblar, M.G., 1991. The Balderton Sand and Gravel: pre-Ipswichian
1501 cold stage fluvial deposits near Lincoln, England. *Journal of Quaternary Science*
1502 **6**, 117–13.
- 1503 Breeze, P.S., Drake, N.A., Groucutt, H.S., Parton, A., Jennings, R.P., White, T.S.,
1504 Clark-Balzan, L., Shipton, C., Scerri, E.M.L., Stimpson, C.M., Al-Omari, A.,
1505 Petraglia, M.D., 2015. Remote sensing and GIS techniques for reconstructing
1506 Arabian palaeohydrology and identifying archaeological sites. *Quaternary*
1507 *International* **382**, 98–119.
- 1508 Breeze, P.S., Groucutt, H.S., Drake, N.A., White, T.S., Jennings, R.P., Petraglia,
1509 M.P., 2016. Palaeohydrological corridors for hominin dispersals in the Middle
1510 East ~250–70,000 years ago. *Quaternary Science Reviews* **144**, 155–185.
- 1511 Briant, R.M., Kilfeather, A.A., Parfitt, S., Penkman, K.E.H., Preece, R.C., Roe, H.M.,
1512 Schwenninger, J.-L., Wenban-Smith, F.F., Whittaker, J.E., 2012. Integrated
1513 chronological control on an archaeologically significant Pleistocene river terrace
1514 sequence: the Thames-Medway, eastern Essex, England. *Proceedings of the*
1515 *Geologists' Association* **123**, 87–108.
- 1516 Bridgland, D.R., 1994. *Quaternary of the Thames*. Geological Conservation Review
1517 Series, Vol. 7. Chapman and Hall, London.
- 1518 Bridgland, D.R., 2000. River terrace systems in north-west Europe: an archive of
1519 environmental change, uplift and early human occupation. *Quaternary Science*
1520 *Reviews* **19**, 1293–1303.
- 1521 Bridgland, D.R., 2006. The Middle and Upper Pleistocene sequence in the Lower
1522 Thames: a record of Milankovitch climatic fluctuation and early human
1523 occupation of southern Britain. *Proceedings of the Geologists' Association* **117**,
1524 281–305.
- 1525 Bridgland, D.R., Schreve, D.C., 2001. River terrace formation in synchrony with
1526 long-term climatic fluctuation: examples from SE England. In: Maddy, D.,
1527 Macklin, M., Woodward, J. (Eds), *River Basin Sediments Systems: Archives of*
1528 *Environmental Change*. Balkema, Rotterdam, 229–248.
- 1529 Bridgland, D.R., Schreve, D.C., 2009. Implications of new Quaternary uplift models
1530 for correlation between the Middle and Upper Thames terrace sequences, UK.
1531 *Global and Planetary Change* **68**, 346–356.
- 1532 Bridgland, D.R., Westaway, R., 2007. Climatically controlled river terrace staircases:
1533 a worldwide Quaternary phenomenon. *Geomorphology* **98**, 285–315.
- 1534 Bridgland, D.R., Westaway, R., 2008a. Preservation patterns of Late Cenozoic fluvial
1535 deposits and their implications: results from IGCP 449. *Quaternary International*
1536 **189**, 5–38.
- 1537 Bridgland, D.R., Westaway, R., 2008b. Climatically controlled river terrace
1538 staircases: a worldwide Quaternary phenomenon. *Geomorphology* **98**, 285–315.

- 1539 Bridgland, D.R., Westaway, R., 2014. Quaternary fluvial archives and landscape
1540 evolution: a global synthesis. *Proceedings of the Geologists' Association* **125**,
1541 600–629
- 1542 Bridgland, D.R. and White, M.J., 2014. Fluvial archives as a framework for the
1543 Lower and Middle Palaeolithic: patterns of British artefact distribution and
1544 potential chronological implications. *Boreas* **43**, 543–555.
- 1545 Bridgland, D.R. and White, M.J., 2015. Chronological variations in handaxes:
1546 patterns detected from fluvial archives in north- west Europe. *Journal of*
1547 *Quaternary Science* **30**, 623–638.
- 1548 Bridgland, D.R., Preece, R.C., Roe, H.M., Tipping, R.M., Coope, G.R., Field, M.H.,
1549 Robinson, J.E., Schreve, D.C., Crowe, K., 2001. Middle Pleistocene interglacial
1550 deposits at Barling, Essex, UK: evidence for a longer chronology for the Thames
1551 terrace sequence. *Journal of Quaternary Science* **16**, 813–840.
- 1552 Bridgland, D.R., Philip, G., Westaway, R., White, M., 2003. A long Quaternary
1553 terrace sequence in the Orontes River valley, Syria: a record of uplift and of
1554 human occupation. *Current Science* **84**, 1080–1089.
- 1555 Bridgland, D.R., Maddy, D., Bates, M., 2004. River terrace sequences: templates for
1556 Quaternary geochronology and marine-terrestrial correlation. *Journal of*
1557 *Quaternary Science* **19**, 203–218.
- 1558 Bridgland, D.R., Antoine, P., Limondin-Lozouet, N., Santisteban, J.I., Westaway, R.,
1559 White, M.J., 2006. The Palaeolithic occupation of Europe as revealed by
1560 evidence from the rivers: data from IGCP 449. *Journal of Quaternary Science* **21**,
1561 437–455.
- 1562 Bridgland, D.R., Field, M.H., Holmes, J.A., McNabb, J., Preece, R.C., Selby, I.,
1563 Wymer, J.J., Boreham, S., Irving, B., Parfitt, S.A., Stuart, A.J., 1999. Middle
1564 Pleistocene interglacial Thames- Medway deposits at Clacton-on-Sea, England:
1565 reconsideration of the biostratigraphical and environmental context of the type
1566 Clactonian Palaeolithic industry. *Quaternary Science Reviews* **18**, 109–146.
- 1567 Bridgland, D.R., Keen, D., Westaway, R., 2007. Global correlation of Late Cenozoic
1568 fluvial deposits: a synthesis of data from IGCP 449. *Quaternary Science Reviews*
1569 **26**, 2694–2700.
- 1570 Bridgland, D.R., Westaway, R., Daoud, M., Yassminh, R., Abou Romieh, M., 2008.
1571 River Terraces of the Nahr el Kebir, NW Syria, and their *Palaeolithic record*.
1572 *Council for British Research in the Levant Bulletin* **3**, 36– 41.
- 1573 Bridgland, D.R., Westaway, R., Romieh, M.A., Candy, I., Daoud, M., Demir, T.,
1574 Galiatsatos, N., Schreve, D.C., Seyrek, A., Shaw, A.D., White, T.S., Whittaker,
1575 J., 2012. The River Orontes in Syria and Turkey: Downstream variation of fluvial
1576 archives in different crustal blocks. *Geomorphology* **165-166**, 25–49.
- 1577 Bridgland, D.R., Harding, P., Allen, P., Candy, I., Cherry, C., George, W., Horne, D.,
1578 Keen, D.H., Penkman, K.E.H., Preece, R.C., Rhodes, E.J., Scaife, R., Schreve,
1579 D.C., Schwenninger, J.-L., Slipper, I., Ward, G., White, M.J., White, T.S.,
1580 Whittaker, J.E., 2013. An enhanced record of MIS 9 geoarchaeology: data from
1581 construction of the High Speed 1 (London – Channel Tunnel) rail-link and other
1582 recent investigations at Purfleet, Essex, UK. *Proceedings of the Geologists'*
1583 *Association* **124**, 417–476.

- 1584 Bridgland, D.R., Howard, A.J., White, M.J., White, T.S., 2014. *Quaternary of the*
1585 *Trent*. Oxbow Books, Oxford.
- 1586 Bridgland, D.R., Howard, A.J., White, M.J., White, T.S., Westaway, R., 2015. New
1587 insight into the Quaternary evolution of the River Trent, UK. *Proceedings of the*
1588 *Geologists' Association* **126**, 466–479.
- 1589 Bridgland, D.R., Demir, T., Seyrek, A., Daoud, M., Abou Romieh, M., Westaway, R.,
1590 2017 (this issue). Drivers of river-terrace formation in the NE Mediterranean
1591 region: evidence from Syria and Turkey. *Quaternary Science Reviews*.
- 1592 Caloi, L., Palombo, M.R., Zarlenga, F., 1998. Late-Middle Pleistocene mammal
1593 faunas of Latium (central Italy): stratigraphy and environment. *Quaternary*
1594 *International* **47**, 77–86.
- 1595 Candy, I., Schreve, D.C., 2007. Land–sea correlation of Middle Pleistocene temperate
1596 sub-stages using high-precision uranium-series dating of tufa deposits from
1597 southern England. *Quaternary Science Reviews* **26**, 1223–1235.
- 1598 Candy, I., Black, S., Sellwood, B.W., 2004. Quantifying timescales of pedogenic
1599 calcrete formation using U-series disequilibria. *Sedimentary Geology* **170**, 177–
1600 187.
- 1601 Candy, I., Black, S., Sellwood, B.W., 2005. U-series isochron dating of immature and
1602 mature calcretes as a basis for constructing Quaternary landform chronologies;
1603 examples from the Sorbas basin, southeast Spain. *Quaternary Research* **64**, 100–
1604 111.
- 1605 Candy, I., Coope, G.R., Lee, J.R., Parfitt, S.A., Preece, R.C., Rose, J., Schreve, D.C.,
1606 2010. Pronounced warmth during early Middle Pleistocene interglacials:
1607 investigating the mid-Bruhnes Event in the British terrestrial sequence. *Earth-*
1608 *Science Reviews* **103**, 183–196.
- 1609 Candy, I., Schreve, D.C., Sherriff, J., Tye, G.J., 2014. Marine Isotope Stage 11:
1610 palaeoclimates, palaeoenvironments and its role as an analogue for the current
1611 interglacial. *Earth-Science Reviews* **128**, 18–51.
- 1612 Candy, I., Schreve, D.C., White, T.S., 2015. MIS 13–12 in Britain and the North
1613 Atlantic: understanding the palaeoclimatic context of the earliest Acheulean.
1614 *Journal of Quaternary Science* **30**, 593–609.
- 1615 Candy, I., White, T.S., Elias, S., 2016 (2016). How warm was Britain during the Last
1616 Interglacial? A critical review of Ipswichian (MIS 5e) palaeotemperature
1617 reconstructions. *Journal of Quaternary Science* **31**, 857–868.
- 1618 Carlini, A.A., Zurita, A.E., Aguilera, O.A., 2008. North American glyptodontines
1619 (Xenarthra, Mammalia) in the Upper Pleistocene of northern South America.
1620 *Paläontologische Zeitschrift* **82**, 125–138.
- 1621 Chauhan, P.R., 2008. Large mammal fossil occurrences and associated archaeological
1622 evidence in Pleistocene contexts of peninsular India and Sri Lanka. *Quaternary*
1623 *International* **192**, 20–42.
- 1624 Chauhan, P.R., Bridgland, D.R., Moncel, M.-H., Antoine, P., Bahain, J.-J., Briant, R.,
1625 Cunha, P., Despriée, J., Limondin-Lozouet, N., Locht, J.-L., Martines, A.,
1626 Schreve, D.C., Shaw, A.D., Voinchet, P., Westaway, R., White, M.J., White,
1627 T.S., 2017 (this issue). Fluvial deposits as an archive of early human activity:

- 1628 progress during the 20 years of the Fluvial Archives Group. *Quaternary Science*
1629 *Reviews*.
- 1630 Chlachula, J., 2010. Environmental context of Pleistocene peopling of the Central
1631 Urals. *Quaternary International* **220**, 47–63.
- 1632 Cione, A.L., Tonni, E.P., 1999. Biostratigraphy and chronological scale of upper-most
1633 Cenozoic in the Pampean area, Argentina. *Quaternary of South America and*
1634 *Antarctic Peninsula* **12**, 23–51.
- 1635 Cione, A.L., Tonni, E.L., 2001 Correlation of Pliocene to Holocene southern South
1636 American and European vertebrate-bearing units. *Bolletino della Società*
1637 *Paleontologica Italiana* **40**, 167–173.
- 1638 Cione, A., Tonni, E.P., 2005. Bioestratigrafía basada en mamíferos del Cenozoico
1639 Superior de la provincia de Buenos Aires, Argentina. In: de Barrio, R.E.,
1640 Etcheverry, R.O., Caballé, M.F., Lambías, E. (Eds), *Relatorio, Geología y*
1641 *Recursos Minerales de la Provincia de Buenos Aires, Congreso Geológico*
1642 *Argentino, 16th, La Plata*. Asociación Geológica Argentina, Buenos Aires, 183–
1643 200.
- 1644 Cliquet, D., Lautridou, J.-P., Antoine, P., Lamotte, A., Leroyer, M., Limondin-
1645 Lozouet, N., Mercier, N., 2009. La séquence loessique de Saint-Pierre-lès-Elbeuf
1646 (Normandie, France) : nouvelles données archéologiques, géochronologiques et
1647 paléontologiques. *Quaternaire* **20**, 321–343.
- 1648 Cohen, T.J., Nanson, G.C., Jansen, J.D., Jones, B.G., Jacobs, Z., Larsen, J.R., May,
1649 J.H., Treble, P., Price, D.M., Smith, A.M., 2012. Late Quaternary mega-lakes fed
1650 by the northern and southern river systems of central Australia: varying moisture
1651 sources and increased continental aridity. *Palaeogeography, Palaeoclimatology,*
1652 *Palaeoecology* **356**, 89–108.
- 1653 Cohen, T.J., Nanson, G.C., Larsen, J.R., Jones, B.G., Price, D.M., Coleman, M.,
1654 Pietsch, T.J., 2010. Late Quaternary aeolian and fluvial interactions on the
1655 Cooper Creek Fan and the association between linear and source-bordering
1656 dunes, Strzelecki Desert, Australia. *Quaternary Science Reviews* **29**, 455–471.
- 1657 Coltorti, M., Abbazzi, L., Ferretti, M.P., Iacumin, P., Paredes Rios, F., Pellegrini, M.,
1658 Pieruccini, P., Rustioni, M., Tito, G., Rook, L., 2007. Last Glacial mammals in
1659 South America: a new scenario from the Tarija Basin (Bolivia).
1660 *Naturwissenschaften* **94**, 288–299.
- 1661 Cooke, H.B.S., 1949. Fossil mammals of the Vaal River deposits. *Memoirs of the*
1662 *Geological Survey of South Africa* **35**, 1–117.
- 1663 Coope, G.R., 2001. Biostratigraphical distinction of interglacial coleopteran
1664 assemblages from southern Britain attributed to Oxygen Isotope Stages 5e and 7.
1665 *Quaternary Science Reviews* **20**, 1717–1722.
- 1666 Coope, G.R., 2010. Coleopteran faunas as indicators of interglacial climates in central
1667 and southern England. *Quaternary Science Reviews* **29**, 1507–1514.
- 1668 Cooper, A., Turney, C., Hughen, K.A., Brook, B.W., McDonald, H.G., Bradshaw,
1669 C.J., 2015. Abrupt warming events drove Late Pleistocene Holarctic megafaunal
1670 turnover. *Science* **349**, 602–606.

- 1671 Cordier, S., Adamson, K., Delmas, M., Calvet, M., Harmand, D., 2017 (this issue).
 1672 Quaternary fluvial response to climate forcing in glacially influenced river
 1673 systems. *Quaternary Science Reviews*.
- 1674 G. Cuenca-Bescós, J. Rofes, J.M. López-García, H.A. Blain, R.J. De Marfá, M.A.
 1675 Galindo-Pellicena, M.L. Bennásar-Serra, M. Melero-Rubio, J.L. Arsuaga, J.M.
 1676 Bermúdez de Castro, E. Carbonell. Biochronology of Spanish Quaternary small
 1677 vertebrate faunas. *Quaternary International* **212**, 109–119
- 1678 Cunha, P.P., Martins, A.A., Huot, S., Murray, A.S., Raposo, L., 2008. Dating the Tejo
 1679 River lower terraces in the Ródão area (Portugal) to assess the role of tectonics
 1680 and uplift. *Geomorphology* **102**, 43–54.
- 1681 Dabkowski, J., Antoine, P., Limondin-Lozouet, N., Chaussé, C., Carbonell, P., 2010.
 1682 Les microfaciès du tuf calcaire éémien (SIM 5e) de Caours (Somme, France) :
 1683 éléments d'analyse paléoécologique du dernier interglaciaire. *Quaternaire* **21**,
 1684 127–137.
- 1685 Dabkowski, J., 2014. High potential of calcareous tufas for integrative
 1686 multidisciplinary studies and prospects for archaeology in Europe. *Journal of*
 1687 *Archaeological Science* **52**, 72–83.
- 1688 Dabowski, J., Limondin-Lozouet, N., Antoine, P., Marca-Bell, A., Andrews, J., 2011.
 1689 Enregistrement des variations climatiques au cours des interglaciaires d'après
 1690 l'étude des isotopes stables de la calcite des tufs calcaires pleistocènes du nord de
 1691 la France : exemples des séquences de Caours (SIM 5e ; Somme) et de La Celle-
 1692 sur-Seine (SIM). *Quaternaire* **22**, 275–285.
- 1693 Dabkowski, J., Limondin-Lozouet, N., Antoine, P. et al. 2012. Climatic variations in
 1694 MIS 11 recorded by stable isotopes and trace elements in a French tufa (La Celle,
 1695 Seine valley). *Journal of Quaternary Science* **27**, 790–799.
- 1696 Dabkowski, J., Royle, S.H., Antoine, P., Marca-Bell, A., Andrews, J.E., 2015. High-
 1697 resolution $\delta^{18}\text{O}$ seasonality record in a French Eemian tufa stromatolite (Caours,
 1698 Somme Basin). *Palaeogeography, Palaeoclimatology, Palaeoecology* **438**, 277–
 1699 284.
- 1700 Danukalova, G.A., Yakovlev, A.G., Morozova, E.M., Alimbekova, L.I., 2007.
 1701 Biostratigraphy of the Late Middle Pleistocene (Middle Neopleistocene) of the
 1702 southern Urals region. *Quaternary International* **160**, 17–29.
- 1703 Danukalova, G., Yakovlev, A., Alimbekova, L., Yakovleva, T., Morozova, E.,
 1704 Ereemeev, A., Kosintsev, P., 2008. Biostratigraphy of the Upper Pleistocene
 1705 (Upper Neopleistocene) – Holocene deposits of the Lemeza River valley of the
 1706 Southern Urals region (Russia). *Quaternary International* **190**, 38–57.
- 1707 Danukalova, G., Yakovlev, A., Kosintsev, P., Agadjanian, A., Alimbekova, L.,
 1708 Ereemeev, A., Morozova, E., 2009. Quaternary fauna and flora of the Southern
 1709 Urals region (Bashkortostan Republic). *Quaternary international* **201**, 13–24.
- 1710 Danukalova, G., Yakovlev, A., Osipova, E., Yakovleva, T., Kosintsev, P., 2011.
 1711 Biostratigraphy of the Upper Pleistocene (Upper Neopleistocene) – Holocene
 1712 deposits of the latitude current of the Belaya river valley of the Southern Urals
 1713 region (Russia). *Quaternary International* **231**, 23–43.

- 1714 Danukalova, G., Yakovlev, A., Osipova, E., Kurmanov, R., van Kolfschoten, T.,
1715 2016. Biostratigraphy of the early Middle Pleistocene of the Southern Fore-Urals.
1716 *Quaternary International* **420**, 115–135.
- 1717 Delagnes, A., Tribolo, C., Bertran, P., Brenet, M., Crassard, R., Jaubert, J., Khalidi,
1718 L., Mercier, N., Nomade, S., Peigné, S., Sitzia, L., Tournepiche, J., Al-halibi, M.,
1719 Al-mosabi, A., Macchiarelli, R., 2012. Inland human settlement in southern
1720 Arabia 55,000 years ago. New evidence from the Wadi Surdud Middle
1721 Paleolithic site complex, western Yemen. *Journal of Human Evolution* **63**, 452–
1722 474.
- 1723 Delagnes, A., Crassard, R., Bertran, P., Sitzia, L., 2013. Cultural and human dynamics
1724 in southern Arabia at the end of the Middle Paleolithic. *Quaternary International*
1725 **300**, 234–243.
- 1726 Demir, T., Westaway, R., Seyrek, A., Bridgland, D., 2007. Terrace staircases of the
1727 River Euphrates in southeast Turkey, northern Syria and western Iraq: evidence
1728 for regional surface uplift. *Quaternary Science Reviews* **26**, 2844–2863.
- 1729 Demir, T., Seyrek, A., Westaway, R., Bridgland, D., Beck, A., 2008. Late Cenozoic
1730 surface uplift revealed by incision by the River Euphrates at Birecik, southeast
1731 Turkey. *Quaternary International* **186**, 132–163.
- 1732 Demir, T., Seyrek, A., Westaway, R., Guillou, H., Scaillet, S., Beck, A., Bridgland,
1733 D.R., 2012. Late Cenozoic regional uplift and localised crustal deformation
1734 within the northern Arabian Platform in southeast Turkey: investigation of the
1735 Euphrates terrace staircase using multidisciplinary techniques. *Geomorphology*
1736 **165-166**, 7–24.
- 1737 Dennell, R.W., 2013. The Nihewan Basin of North China in the Early Pleistocene:
1738 continuous and flourishing, or discontinuous, infrequent and ephemeral
1739 occupation? *Quaternary International* **295**, 223–236.
- 1740 Dennell, R.W., Coard, R., Turner, A., 2008. Predators and scavengers in Early
1741 Pleistocene southern Asia. *Quaternary International* **192**, 78–88.
- 1742 Deschamps, C.M., 2005. Late Cenozoic mammal bio-chronostratigraphy in
1743 southwestern Buenos Aires Province, Argentina. *Ameghiniana* **42**, 733–750.
- 1744 Di Stefano, G., Petronio, C., Sardella, R., 1998. Biochronology of the Pleistocene
1745 mammal Faunas from Rome urban area. *Il Quaternario* **11**, 191–199.
- 1746 Dortch, J., Cupper, M., Grün, R., Harpley, B., Lee, K. and Field, J., 2016. The timing
1747 and cause of megafauna mass deaths at Lancefield Swamp, south-eastern
1748 Australia. *Quaternary Science Reviews* **145**, 161–182.
- 1749 Drake, N.A., Blench, R.M., Armitage, S.J., Bristow, C.S., White, K.H., 2011. Ancient
1750 watercourses and biogeography of the Sahara explain the peopling of the desert.
1751 *Proceedings of the National Academy of Sciences* **108**, 458–62.
- 1752 Dykan, N., 2003. The biostratigraphic division of the Neopleistocene deposits of
1753 Ukraine according to ostracoda fauna. *Geological Journal* **3**, 114–120.
- 1754 Edgell, H.S., 2006. Evolution of Arabian deserts and their chronology. In: Edgell,
1755 H.S. (Ed.), *Arabian Deserts: Nature, Origin, and Evolution*. Springer, New York,
1756 493–511.

- 1757 Elton, S., O'Regan, H.J., 2014. Macaques at the margins: the biogeography and
1758 extinction of *Macaca sylvanus* in Europe. *Quaternary Science Reviews* **96**, 117–
1759 130.
- 1760 Fauzi, M.R., Ansyori, M.M., Prastiningtyas, D., Intan, M.F.S., Wibowo, U.P.,
1761 Rahmanendra, H., Widiyanto, H. and Simanjuntak, T., 2016. Matar: A forgotten
1762 but promising Pleistocene locality in East Java. *Quaternary International* **416**,
1763 183–192.
- 1764 Garcia, J., 2015. Long-term hominin occupation during the Middle and early Late
1765 Pleistocene: chronostratigraphy of the Ter River basin (Girona, north-eastern
1766 Spain). *Journal of Archaeological Science Reports* **3**, 493–512.
- 1767 Gaudenyi, T., Nenadic, D., Jovanovic, M., Bogicevic, K., 2013. The stratigraphical
1768 importance of *Viviparus boeckhi* Horizon of Serbia. *Quaternary International*
1769 **292**, 101–112.
- 1770 Gaudenyi, T., Nenadic, Stejic, P., Jovanovic, M., Bogicevic, K., 2015a. Stratigraphic
1771 revision of the Quaternary material from the vicinity of Belgrade and the Serbian
1772 segment of the Tisza valley analysed by V.D. Laskarev. *Quaternary International*
1773 **357**, 93–109.
- 1774 Gaudenyi, T., Nenadic, D., Stejic, P., Jovanovic, M., Bogicevic, K., 2015b. The
1775 stratigraphy of the Serbian Pleistocene *Corbicula* beds. *Quaternary International*
1776 **357**, 4–21.
- 1777 Geraards, D., Alemseged, Z., Reed, D., Wynn, J., Roman, D.C., 2004. The
1778 Pleistocene fauna (other than primates) from Asbole, lower Awash Valley,
1779 Ethiopia, and its environmental and biochronological implications. *Geobios* **37**,
1780 697–718.
- 1781 Girotti, O., Capasso Barbato, L., Esu, D., Gliozzi, E., Kotsakis, T., Martinetto, E.,
1782 Petronio, C., Sardella, R., Squazzini, E., 2003. The section of Torre di Picchio
1783 (Terni, Umbria, Central Italy): a Villafranchian site rich in Vertebrates, Molluscs,
1784 Ostracods and Plants. *Rivista Italiana Paleontologia e Stratigrafia* **109**, 77–98.
- 1785 Gleed-Owen, C.P., 1988. Quaternary herpetofaunas of the British Isles: taxonomic
1786 descriptions, palaeoenvironmental reconstructions, and biostratigraphic
1787 implications. Unpublished Ph.D. thesis, Coventry University.
- 1788 Gleed-Owen, C.P., 1999. The palaeoclimatic and biostratigraphic significance of
1789 herpetofaunal remains from the British Quaternary. In: Andrews, P., Banham, P.
1790 (Eds), *Late Cenozoic Environments and Hominid Evolution: a tribute to Bill*
1791 *Bishop*. Geological Society, London, 201–215.
- 1792 Gliozzi, E., Abbazzi, L., Argenti, P., Azzaroli, A., Caloi, L., Capasso Barbato, L., Di
1793 Stefano, G., Esu, D., Ficarelli, G., Girotti, O., Kotsakis, T., Masini, F., Mazza,
1794 P., Mezzabotta, C., Palombo, M.R., Petronio, C., Rook, L., Sala, B., Sardella, R.,
1795 Zanolida, E., Torre, D., 1997. Biochronology of selected mammals, molluscs and
1796 ostracods from the Middle Pliocene to the Late Pleistocene in Italy: the state of
1797 the art. *Rivista Italiana di Paleontologia e Stratigrafia* **103**, 369–388.
- 1798 Griffiths, H.I., 2001. Ostracod evolution and extinction – its biostratigraphic value in
1799 the European Quaternary. *Quaternary Science Reviews* **20**, 1743–1751.
- 1800 Green, C.P., Branch, N.P., Coope, G.R., Field, M.H., Keen, D.H., Wells, J.M.,
1801 Schwenninger, J.-L., Preece, R.C., Schreve, D.C., Canti, G., Gleed-Owen, C.P.,

- 1802 2006. Marine Isotope Stage 9 environments of fluvial deposits at Hackney, North
1803 London, UK. *Quaternary Science Reviews* **25**, 89–113.
- 1804 Groucutt, H.S., Petraglia, M.D., 2012. The prehistory of the Arabian Peninsula:
1805 deserts, dispersals, and demography. *Evolutionary Anthropology* **21**, 113–25
- 1806 Groucutt, H.S., White, T.S., Clark-Balzan, L., Parton, A., Crassard, R., Shipton, C.,
1807 Jennings, R.P., Parker, A.G., Breeze, P.S., Scerri, E.M.L., Alsharekh, A.,
1808 Petraglia, M.D., 2015. Human occupation of the Arabian Empty Quarter during
1809 MIS 5: evidence from Mundafan Al-Buhayrah, Saudi Arabia. *Quaternary*
1810 *Science Reviews* **119**, 116–135
- 1811 Habeck-Fardy, A., Nanson, G.C., 2014. Environmental character and history of the
1812 Lake Eyre Basin, one seventh of the Australian continent. *Earth-Science Reviews*
1813 **132**, 39–66.
- 1814 Haynes, G., 2009. *American Megafaunal Extinctions at the End of the Pleistocene*.
1815 Springer, New York.
- 1816 Helgren, D.M., 1977. Geological context of the Vaal River faunas. *South African*
1817 *Journal of Science* **73**, 303–307.
- 1818 Hérissou, D., Loch, J.-L., Vallin, L., Deschodt, L., Antoine, P., Auguste, P.,
1819 Limondin-Lozouet, N., Lefèvre, S., Hulin, G., Masson, B., Ghaleb, B., 2015.
1820 Neandertal's presence during the Eemian interglacial in North-West Europe: a
1821 new site at Waziers (Northern France). *Proceedings of the European Society for*
1822 *the study of Human Evolution* **4**, 115.
- 1823 Hilbert, Y.H., White, T.S., Parton, A., Clark-Balzan, L., Crassard, R., Groucutt, H.S.,
1824 Jennings, R.P., Breeze, P., Parker, A., Shipton, C., Al-Omari, A., Alsharekh,
1825 A.M., Petraglia, M.D., 2014. Epipalaeolithic occupation and palaeoenvironments
1826 of the southern Nefud desert, Saudi Arabia, during the Terminal Pleistocene and
1827 Early Holocene. *Journal of Archaeological Science* **50**, 460–474.
- 1828 Holm, D., 1960. Desert Geomorphology in the Arabian Peninsula. *Science* **132**, 1369–
1829 1379.
- 1830 Holman, J.A., 1998. *Pleistocene amphibians and reptiles in Britain and Europe*.
1831 Oxford University Press, Oxford.
- 1832 Holyoak, D.T., Preece, R.C., 1985. Late Pleistocene interglacial deposits at
1833 Tattershall, Lincolnshire. *Philosophical Transactions of the Royal Society of*
1834 *London B* **331**, 193–236.
- 1835 Horne, D.J., 2007. A mutual temperature range method for Quaternary palaeoclimatic
1836 analysis using European nonmarine Ostracoda. *Quaternary Science Reviews* **26**,
1837 1398–1415.
- 1838 Horne, D.J., Curry, B.B., Mesquita-Joanes, F., 2012. Mutual climatic range methods
1839 for Quaternary ostracods. *Developments in Quaternary Science* **17**, 65–84.
- 1840 Howard, A.J., Keen, D.H., Hollin, J.T., 1999. Amino acid dating of a molluscan fauna
1841 from Bassingham Fen, Lincolnshire: implications for the chronology of the Trent
1842 Terraces. *Proceedings of the Geologists' Association* **110**, 233–239.
- 1843 Howard, A.J., Bridgland, D., Knight, D., McNabb, J., Rose, J., Schreve, D.,
1844 Westaway, R., White, M.J. and White, T.S., 2007. The British Pleistocene fluvial
1845 archive: East Midlands drainage evolution and human occupation in the context

- 1846 of the British and NW European record. *Quaternary Science Reviews* **26**, 2724–
1847 2737.
- 1848 Jado, R.A., Zötl, J., 1984. *The Quaternary Period in Saudi Arabia, Vol.2.*
1849 *Sedimentological, Hydrogeological, Hydrochemical, Geomorphological,*
1850 *Geochronological and Climatological Investigations in Western Saudi Arabia.*
1851 Springer-Verlag, New York.
- 1852 Jaeger, J.J., Wesselman, H.B., 1976. Fossil remains of micromammals from the Omo
1853 Group deposits. In: Coppens, Y., Howells, F.C., Isaac, G.L., Leakey, R.E.F. (Eds)
1854 *Earliest Man and Environments in the Lake Rudolph Basin: Stratigraphy,*
1855 *Paleoecology, and Evolution.* University of Chicago Press, London, 351–360.
- 1856 Jennings, R.P., Singarayer, J., Stone, E.J., Krebs-kanzow, U., Khon, V., Nisancioglu,
1857 K.H., Pfeiffer, M., Zhang, X., Parker, A., Parton, A., Groucutt, H.S., White, T.S.,
1858 Drake, N.A., Petraglia, M.D., 2015a. The greening of Arabia: multiple
1859 opportunities for human occupation of the Arabian Peninsula during the Late
1860 Pleistocene inferred from an ensemble of climate model simulations. *Quaternary*
1861 *International* **382**, 181–199.
- 1862 Jennings R.P., Parton, A., Clark-Balzan, L., White, T.S., Groucutt, H.S., Breeze, P.,
1863 Parker, A., Drake, N.A., Petraglia, M.P., 2017. Human occupation of the northern
1864 Arabian interior during early Marine Isotope Stage 3. *Journal of Quaternary*
1865 *Science* **31**, 953–966.
- 1866 Johnson, C.N., 2017. Fire, people and ecosystem change in Pleistocene
1867 Australia. *Australian Journal of Botany* **64**, 643–651.
- 1868 Johnson, C.N., Alroy, J., Beeton, N.J., Bird, M.I., Brook, B.W., Cooper, A., Gillespie,
1869 R., Herrando-Pérez, S., Jacobs, Z., Miller, G.H., Prideaux, G.J., 2016. What
1870 caused extinction of the Pleistocene megafauna of Sahul? *Proceedings of the*
1871 *Royal Society of London B: Biological Sciences* **283**, 20152399.
- 1872 Jones, P.F., Stanley, M.F., 1974. Ipswichian mammalian fauna from the Beeston
1873 Terrace at Boulton Moor, near Derby. *Geological Magazine* **111**, 515–520.
- 1874 Jones, P.F., Stanley, M.F., 1975. Description of Hippopotamus and other mammalian
1875 remains from the Allenton Terrace of the Lower Derwent Valley. *The Mercian*
1876 *Geologist* **5**, 259–272.
- 1877 Kahlke, R.D., García, N., Kostopoulos, D.S., Lacombe, F., Lister, A.M., Mazza, P.P.,
1878 Spassov, N., Titov, V.V., 2011. Western Palaeartic palaeoenvironmental
1879 conditions during the Early and early Middle Pleistocene inferred from large
1880 mammal communities, and implications for hominin dispersal in Europe.
1881 *Quaternary Science Reviews* **30**, 1368–1395.
- 1882 Keen, D.H., 1990. Significance of the record provided by Pleistocene fluvial deposits
1883 and their included molluscan faunas for palaeoenvironmental reconstruction and
1884 stratigraphy: case study from the English Midlands. *Palaeogeography,*
1885 *Palaeoclimatology, Palaeoecology* **80**, 25–34.
- 1886 Kemp, J., Rhodes, E.J. 2010. Episodic fluvial activity of inland rivers in southeastern
1887 Australia: palaeochannel systems and terraces of the Lachlan River. *Quaternary*
1888 *Science Reviews* **29**, 732–752.

- 1889 Kemp, J. and Spooner, N.A., 2007. Evidence for regionally wet conditions before the
1890 LGM in southeast Australia: OSL ages from a large palaeochannel in the Lachlan
1891 Valley. *Journal of Quaternary Science* **22**, 423–427.
- 1892 Kennard, A.S., 1942a. Some Pleistocene problems discussed (discussion appended to
1893 Bull, A.J., 1942. *Proceedings of the Geologists' Association* **53**, 24–25.
- 1894 Kennard, A.S., 1942b. Faunas of the High Terrace at Swanscombe. *Proceedings of*
1895 *the Geologists' Association* **53**, 105.
- 1896 Kerney, M.P., 1971. Interglacial deposits at Barnfield Pit, Swanscombe, and their
1897 molluscan fauna. *Journal of the Geological Society of London* **127**, 69–86.
- 1898 Kershaw, A.P., Bretherton, S.C., van der Kaars, S., 2007. A complete pollen record of
1899 the last 230 ka from Lynch's Crater, north-eastern Australia. *Palaeogeography,*
1900 *Palaeoclimatology, Palaeoecology* **251**, 23–45.
- 1901 Klein, R.G., 1983. The Stone Age prehistory of southern Africa. *Annual Review of*
1902 *Anthropology* **12**, 25–48.
- 1903 Koch, P.L., Barnosky, A.D., 2006. Late Quaternary extinctions: state of the debate.
1904 *Annual Review of Ecology, Evolution, and Systematics* (2006), 215–250.
- 1905 Kock, S. Kramers, J.D., Preusser, F., Wetzel, A., 2009. Dating of Late Pleistocene
1906 terrace deposits of the River Rhine using Uranium series and luminescence
1907 methods: potential and limitations. *Quaternary Geochronology* **4**, 363–373.
- 1908 Kondrashov, P.E., 2007. New gastropod species from the Pleistocene of the Upper
1909 Don basin. *Paleontological Journal* **41**, 513–519.
- 1910 Kotlia, B.S., Joshi, M., 2011. Taphonomy of Late Pleistocene micromammalian fauna
1911 of Narmada Valley, Central India. *Palaeoworld* **20**, 84–91.
- 1912 Koutsodendris, A., Müller, U.C., Pross, J., Brauer, A., Kotthoff, U., Lotter, A.F.,
1913 2010. Vegetation dynamics and climate variability during the Holsteinian
1914 interglacial based on a pollen record from Dethlingen (northern Germany).
1915 *Quaternary Science Reviews* **29**, 3298–3307.
- 1916 Koutsodendris, A., Brauer, A., Pälike, H., Müller, U.C., Dulski, P., Lotter, A.F.,
1917 Pross, J., 2011. Sub-decadal-to decadal-scale climate cyclicity during the
1918 Holsteinian interglacial (MIS 11) evidenced in annually laminated sediments.
1919 *Climate of the Past* **7**, 987–999.
- 1920 Kuzmina, S., 2014. New approach to the Quaternary studies: QINSIB - the database
1921 of Siberian fossil insects. *Quaternary International* **341**, 283–293.
- 1922 Langford, H.E., Briant, R.M., 2004. *Nene Valley: Field Guide*. Quaternary Research
1923 Association, London.
- 1924 Langford, H.E., Boreham, S., Briant, R.M., Coope, G.R., Horne, D.J., Schreve, D.C.,
1925 Whittaker, J.E., Whitehouse, N.J., 2014. Middle to Late Pleistocene
1926 palaeoecological reconstructions and palaeotemperature estimates for cold/cool
1927 stage deposits at Whittlesey, eastern England. *Quaternary International* **341**, 6–
1928 26.
- 1929 Langford, H.E., Boreham, S., Coope, G.R., Fletcher, W., Horne, D.J., Keen, D.H.,
1930 Mighall, T., Penkman, K.E.H., Schreve, D.C. and Whittaker, J.E., 2014.
1931 Palaeoecology of a late MIS 7 interglacial deposit from eastern England.
1932 *Quaternary International* **341**, 27–45.

- 1933 Limondin-Lozouet, N., 2011. Successions malacologiques à la charnière
1934 Glaciaire/Interglaciaire : du modèle Tardiglaciaire-Holocène aux transitions du
1935 Pléistocène. *Quaternaire* **22**, 211–220.
- 1936 Limondin-Lozouet, N., Gauthier A. Preece, R.C., 2005. Enregistrement des
1937 biocénoses de la première moitié de l'Holocène en contexte tufacé à Saint-
1938 Germain-le-Vasson (Calvados). *Quaternaire* **16**, 265–271.
- 1939 Limondin-Lozouet, N., Antoine, P., 2006. A new *Lyrodiscus* (Mollusca, Gastropoda)
1940 assemblage at Saint-Acheul (Somme Valley): reappraisal of Stage 11
1941 malacofaunas from Northern France. *Boreas* **35**, 622–633.
- 1942 Limondin-Lozouet, N., Antoine, P., Auguste, P., Bahain, J.J., Carbonel, P., Chaussé,
1943 C., Connet, N., Dupéron, J., Dupéron, M., Falguères, C., Freytet, P., Ghaleb, B.,
1944 Jolly-Saad, M.C., Lhomme, V., Lozouet, P., Mercier, N., Pastre, J.F., Voinchet,
1945 P., 2006. Le tuf calcaire de La Celle-sur-Seine (Seine et Marne): nouvelles
1946 données sur un site clé du stade 11 dans le Nord de la France. *Quaternaire* **17**, 5–
1947 29.
- 1948 Limondin-Lozouet, N., Nicoud, E., Antoine, P., Auguste, P., Bahain, J.-J.,
1949 Dabkowski, J., Dupéron, J., Dupéron, M., Falguères, C., Ghaleb, B., Jolly-Saad,
1950 M.-C., Mercier, N., 2010. Oldest evidence of Acheulean occupation in the Upper
1951 Seine valley (France) from an MIS 11 tufa at La Celle. *Quaternary International*
1952 **223-224**, 299–311.
- 1953 Limondin-Lozouet, N., Preece, R.C., 2014. Quaternary perspectives on the diversity
1954 of land snail assemblages from northwestern Europe. *Journal of Molluscan*
1955 *Studies* **80**, 224–237.
- 1956 Limondin-Lozouet, N., Antoine, P., Bahain, J.J., Cliquet, D., Coutard, S., Dabkowski,
1957 J., Ghaleb, B., Locht, J.L., Nicoud, E., Voinchet, P., 2015. North-West European
1958 MIS 11 malacological successions: a framework for the timing of Acheulean
1959 settlements. *Journal of Quaternary Science* **30**, 702–712.
- 1960 Limondin-Lozouet, N., Villa V., Pereira A., Nomade S., Bahain J.J., Stoetzel E.,
1961 Aureli D., Nicoud E., 2017. Middle Pleistocene molluscan fauna from the Valle
1962 Giumentina (Abruzzo, Central Italy): Palaeoenvironmental, biostratigraphical and
1963 biogeographical implications. *Quaternary Science Reviews* **156**, 135–149.
- 1964 Lisiecki, L.E., Raymo, M.E., 2005. A Pliocene-Pleistocene stack of 57 globally
1965 distributed benthic $\delta^{18}\text{O}$ records. *Paleoceanography* **20**, 1–17.
- 1966 Liu, C.R., Yin, G.M., Deng, C.L., Han, F., Song, W.J., 2014. ESR dating of the
1967 Majuangou and Banshan Paleolithic sites in the Nihewan Basin, North China.
1968 *Journal of Human Evolution* **73**, 58–63.
- 1969 Locht, J.L., Coutard, S., Antoine, P., Sellier, N., Ducrocq, T., Paris, C., Guerlin, O.,
1970 Kiefer, D., Defaux, F., Deschodt, L., Limondin-Lozouet, N., 2013. Données
1971 inédites sur le Quaternaire et le Paléolithique du Nord de la France. *Revue*
1972 *archéologique de Picardie* **3/4**, 5–70.
- 1973 Lopes, R.P., 2013. Biostratigraphy of the Pleistocene fossiliferous deposits of the
1974 southern Brazilian coastal area. *Journal of Mammal Evolution* **20**, 69–82.
- 1975 Lopes, R.P., Oliveira, L.C., Figueiredo, A.M.G., Kinoshita, A., Baffa, O., Buchmann,
1976 F.S.C., 2010. ESR dating of Pleistocene mammal teeth and its implications for

- 1977 the biostratigraphy and geological evolution of the coastal plain, Rio Grande do
1978 Sul, southern Brazil. *Quaternary International* **212**, 213–222.
- 1979 Lopes, R.P., Ribeiro, A.M., Dillenburg, S.R., Schultz, C.L., 2013. Late Middle to Late
1980 Pleistocene paleoecology and paleoenvironments in the coastal plain of Rio
1981 Grande do Sul State, southern Brazil, from stable isotopes in fossils of *Toxodon*
1982 and *Stegomastodon*. *Palaeogeography, Palaeoclimatology, Palaeoecology* **369**,
1983 385–394.
- 1984 Lopes, R.P., Kinoshita, A., Baffa, O., Figueiredo, A.M.G., Dillenburg, S.R., Schultz,
1985 C.L., Pereira, J.C., 2014. ESR dating of Pleistocene mammals and marine shells
1986 from the coastal plain of Rio Grande do Sul state, southern Brazil. *Quaternary*
1987 *International* **352**, 124–134.
- 1988 Louys, J., Meijaard, E., 2010. Palaeoecology of Southeast Asian megafauna-bearing
1989 sites from the Pleistocene and a review of environmental changes in the region.
1990 *Journal of Biogeography* **37**, 1432–1449.
- 1991 Louys, J., Curnoe, D., Tong, H., 2007. Characteristics of Pleistocene megafauna
1992 extinctions in Southeast Asia. *Palaeogeography, Palaeoclimatology,*
1993 *Palaeoecology* **243**, 152–173.
- 1994 Maddy, D., Demir, T., Veldkamp, T., Bridgland, D., Stemerink, C., van der Schriek,
1995 T., Schreve, D. 2012. The obliquity-controlled early Pleistocene sequence of the
1996 Gediz River, western Turkey: a revised correlation and chronology. *Journal of the*
1997 *Geological Society of London* **169**, 67–82.
- 1998 Maddy, D., Schreve, D., Demir, T., Veldkamp, A., Wijbrans, J.R., van Gorp, W., van
1999 Hinsbergen, D.J.J., Dekkers, M.J., Scaife, R., Schoorl, J.M., Stemerink, C., van
2000 der Schriek, T. 2015. The earliest securely-dated hominin artefact in Anatolia?
2001 *Quaternary Science Reviews* **109**, 68–75.
- 2002 Maizels, J.K., 1987. Plio-Pleistocene raised channel systems of the western Sharqiya
2003 (Wahiba), Oman. In: Frostick, L., Reid, I. (Eds.) *Desert Sediments: Ancient and*
2004 *Modern*. Geological Society, London, Special Publication No 35, Blackwell
2005 Scientific Publications, Oxford, 31–50.
- 2006 Maizels, J., 1990. Raised channel systems as indicators of palaeohydrologic change: a
2007 case study from Oman. *Palaeogeography, Palaeoclimatology, Palaeoecology* **76**,
2008 241–277.
- 2009 Mania, D., 1995. The earliest occupation of Europe: the Elbe-Saale region
2010 (Germany). In: Roebroeks, W., van Kolfschoten, T. (Eds), *The Earliest*
2011 *Occupation of Europe*. University of Leiden, The Netherlands, 85–101.
- 2012 Mania, D., Altermann, M., 2004. Geology of the Lower Palaeolithic find horizon of
2013 Bilzingsleben (Thuringia) and the anthropogenous aspect. *Hercynia* **38**, 143–184.
- 2014 Markova, A.K., 2006. Likhvin Interglacial small mammal faunas of Eastern Europe.
2015 *Quaternary International* **149**, 67–79.
- 2016 Markova, A.K., 2007. Pleistocene mammal faunas of Eastern Europe. *Quaternary*
2017 *International* **160**, 100–111.
- 2018 Markova, A.K., van Kolfschoten, T., 2012. Middle Pleistocene small mammal faunas
2019 of Eastern and Western Europe: chronology, correlation. *Geography,*
2020 *Environment, Sustainability* **4**, 17–23.

- 2021 Martins, A.A., Cunha, P.P., Buylaert, J.-P., Huot, S., Murray, A.S., Dinis, P., Stokes,
2022 M., 2010. K-Feldspar IRSL dating of a Pleistocene river terrace staircase
2023 sequence of the Lower Tejo River (Portugal, western Iberia). *Quaternary*
2024 *Geochronology* **5**, 176–180.
- 2025 Marra, F., Pandolfi, L., Petronio, C., Di Stefano, G., Gaeta, M., Salari, L., 2014.
2026 Reassessing the sedimentary deposits and vertebrate assemblages from Ponte
2027 Galeria area (Rome, central Italy): An archive for the Middle Pleistocene faunas
2028 of Europe. *Earth-Science Reviews* **139**, 104–122.
- 2029 Marra, F., Pandolfi, L., Petronio, C., Di Stefano, G., Gaeta, M., Salari, L., 2015.
2030 Reply to comment on "Reassessing the sedimentary deposits and vertebrate
2031 assemblages from Ponte Galeria area (Rome, central Italy): an archive for the
2032 Middle Pleistocene faunas of Europe" by Sardella et al. *Earth-Science Reviews*
2033 **147**, 204–205.
- 2034 Marra, F., Florindo, F., Anzidei, M., Sepe, V. 2016. Paleo-surfaces of glacio-
2035 eustatically forced aggradational successions in the coastal area of Rome:
2036 Assessing interplay between tectonics and sea-level during the last ten
2037 interglacials. *Quaternary Science Reviews* **148**, 85–100.
- 2038 Matoshko, A., Gozhik, P., Ivchenko, A., 2002. The fluvial archive of the Middle and
2039 Lower Dnieper (a review). *Geologie en Mijnbouw/ Netherlands Journal of*
2040 *Geoscience* **81**, 339–355.
- 2041 Matoshko, A.V., Gozhik, P., Danukalova, G., 2004. Key Late Cenozoic fluvial
2042 archives in the central and southern part of the East European Plain (a review).
2043 *Proceedings of the Geologists' Association* **115**, 141–173.
- 2044 McLaren, S.J., Al-Juaidi, F., Bateman, M.D., Millington, A.C., 2009. First evidence
2045 for episodic flooding events in the arid interior of central Saudi Arabia over the
2046 last 60 ka. *Journal of Quaternary Science* **24**, 198–207.
- 2047 Megirian, D., Prideaux, G.J., Murray, P.F., Smit, N., 2010. An Australian land
2048 mammal age biochronological scheme. *Paleobiology* **36**, 658–671.
- 2049 Meijer, T., 1988. Fossiele Zoetwatermerieten uit het Nederlandse Kwartair en enkele
2050 opmerkingen over het voorkomen van deze groep in het Kwartair van Noordwest
2051 Europa. *De Kreukel, Jubileumnummer*, 89–108.
- 2052 Meijer, T., Preece, R.C., 1996. Malacological evidence relating to the stratigraphical
2053 position of the Cromerian. In: Turner, C. (Ed.), *The Early Middle Pleistocene in*
2054 *Europe*. Balkema, Rotterdam, 53–82.
- 2055 Meijer, T., Cleveringa, P., 2009. Aminostratigraphy of Middle and Late Pleistocene
2056 deposits in The Netherlands and the southern part of the North Sea Basin. *Global*
2057 *and Planetary Change* **68**, 326–345.
- 2058 Meng, S., Wansa, S., 2005. Lithologie, Stratigraphie und Paläoökologie des
2059 Mittelpleistozäns von Uichteritz im Markkröhlitzer Tal (Lkr. Weißenfels/Sachsen-
2060 Anhalt). *Eiszeitalter und Gegenwart* **55**, 174–214.
- 2061 Meng, S., Wansa, S., 2008. Sediments and processes at the borderline of Saalian
2062 glaciation southwest of Halle (Saale). *Zeitschrift der Deutschen Gesellschaft für*
2063 *Geowissenschaften* **159**, 205–220.

- 2064 Milli, S., Palombo, M.R., Petronio, C., Sardella, R., 2004. The Middle Pleistocene
2065 deposits of the Roman Basin (Latium, Italy): an integrated approach of mammal
2066 biochronology and sequence stratigraphy. *Rivista Italiana di Paleontologia e*
2067 *Stratigrafia* **110**, 559–569.
- 2068 Mishra, S., White, M.J., Beaumont, P., Antoine, P., Bridgland, D.R., Limondin-
2069 Lozouet, N., Santisteban, J.I., Schreve, D.C., Shaw, A.D., Wenban-Smith, F.F.,
2070 Westaway, R.W.C., White, T.S., 2007. Fluvial deposits as an archive of early
2071 human activity. *Quaternary Science Reviews* **26**, 2996–3016.
- 2072 Mishra, S., Gaillard, C., Hertler, C., Moigne, A.M., Simanjuntak, T., 2010. India and
2073 Java: contrasting records, intimate connections. *Quaternary International* **223**,
2074 265–270.
- 2075 Mosquera, M., Saladié, P., Ollé, A., Cáceres, I., Huguet, R., Villalaín, J.J., Carrancho,
2076 A., Bourlès, D., Braucher, R., Vallverdú, J., 2015. Barranc de la Boella
2077 (Catalonia, Spain): an Acheulean elephant butchering site from the European late
2078 Early Pleistocene. *Journal of Quaternary Science* **30**, 651–666.
- 2079 Murphy, B.P., Williamson, G.J., Bowman, D.M.J.S., 2011. Did central Australian
2080 megafaunal extinction coincide with abrupt ecosystem collapse or gradual
2081 climatic change? *Global Ecology and Biogeography* **21**, 142–151.
- 2082 Murton, J.B., Baker, A., Bowen, D.Q., Caseldine, C.J., Coope, G.R., Currant, A.P.,
2083 Evans, J.G., Field, M.H., Green, C.P., Hatton, J., Ito, M., 2001. A late Middle
2084 Pleistocene temperate–periglacial–temperate sequence (Oxygen Isotope Stages
2085 7–5e) near Marsworth, Buckinghamshire, UK. *Quaternary Science Reviews* **20**,
2086 1787–1825.
- 2087 Nanda, A.C., 2002. Upper Siwalik mammalian faunas of India and associated events.
2088 *Journal of Asian Earth Sciences* **21**, 47–58.
- 2089 Nanda, A.C., 2008. Comments on the Pinjor Mammalian Fauna of the Siwalik Group
2090 in relation to the post-Siwalik faunas of Peninsular India and Indo-Gangetic
2091 Plain. *Quaternary International* **192**, 6–13.
- 2092 Pandolfi, L., Marra, F., 2015. Rhinocerotidae (Mammalia, Perissodactyla) from the
2093 chrono-stratigraphically constrained Pleistocene deposits of the urban area of
2094 Rome (Central Italy). *Geobios* **48**, 147–167.
- 2095 Pandolfi, L., Petronio, C., 2011a. *Stephanorhinus etruscus* (Falconer, 1868) from
2096 Pirro Nord (Apricena, Foggia, Southern Italy) with notes on the other late Early
2097 Pleistocene rhinoceros remains of Italy. *Rivista Italiana di Palaeontologia e*
2098 *Stratigrafia* **117**, 173–187.
- 2099 Pandolfi, L. and Petronio, C., 2011b. The small-sized rhinoceroses from the Late
2100 Pleistocene of Apulia (southern Italy). *Rivista Italiana di Paleontologia e*
2101 *Stratigrafia* **117**, 509–520.
- 2102 Pandolfi, L., Gaeta, M., Petronio, C., 2013. The skull of *Stephanorhinus hemitoechus*
2103 (Mammalia, Rhinocerotidae) from the Middle Pleistocene of Campagna Romana
2104 (Rome, central Italy): biochronological and paleobiogeographic implications.
2105 *Bulletin of Geosciences* **88**, 51–62.
- 2106 Parfitt, S.A., Ashton, N.M., Lewis, S.G., Abel, R.L., Coope, G.R., Field, M.H., Gale,
2107 R., Hoare, P.G., Larkin, N.R., Lewis, M.D., Karloukovski, V., Maher, B.A.,
2108 Peglar, S.M., Preece, R.C., Whittaker, J.E., Stringer, C.B. 2010. Early Pleistocene

- 2109 human occupation at the edge of the boreal zone in northwest Europe. *Nature*
2110 **466**, 229–233.
- 2111 Parker, A.G., 2009. Pleistocene climate change in Arabia: developing a framework
2112 for hominin dispersal over the last 350 ka. In: Petraglia, M.D., Rose, J.I. (Eds),
2113 *The Evolution of Human Populations in Arabia: Palaeoenvironments, Prehistory*
2114 *and Genetics*. Springer, New York, 39–49.
- 2115 Parton, A., Farrant, A.R., Leng, M.J., Schwenninger, J., Rose, J.I., Uerpman, H.,
2116 Parker, A.G., 2013. An early MIS 3 pluvial phase in Southeast Arabia: Climatic
2117 and archaeological implications. *Quaternary International* **300**, 62–74.
- 2118 Parton, A., White, T.S., Parker, A.G., Breeze, P.S., Jennings, R., Groucutt, H.S.,
2119 Petraglia, M.D., 2015a. Orbital- scale climate variability in Arabia as a potential
2120 motor for human dispersals. *Quaternary International* **282**, 82–97.
- 2121 Parton, A., Farrant, A.R., Leng, M.J., Telfer, M.W., Groucutt, H.S., Petraglia, M.D.,
2122 Parker, A.G., 2015b. Alluvial fan records from southeast Arabia reveal multiple
2123 windows for human dispersal. *Geology* **43**, 1– 4.
- 2124 Patnaik, R., 2000. Paleoenvironment of *Homo erectus* bearing Pleistocene Narmada
2125 valley deposits of India. *Bulletin of the Oil and Natural Gas Corporation Limited*
2126 **37**, 247–249.
- 2127 Patnaik, R. and Nanda, A.C., 2010. Early Pleistocene mammalian faunas of India and
2128 evidence of connections with other parts of the world. In Fleagle, J.G., Shea, J.J.,
2129 Grine, F.E., Baden, A.L., Leakey, R.E. (Eds), *Out of Africa I: The First Hominin*
2130 *Colonization of Eurasia*. Springer, Netherlands, 129–143.
- 2131 Patnaik, R., Chauhan, P.R., Rao, M.R., Blackwell, B.A.B., Skinner, A.R., Khan, H.S.,
2132 2009. New geochronological, palaeoclimatological and palaeolithic data from the
2133 Narmada Valley hominin locality, central India. *Journal of Human Evolution* **56**,
2134 114–133.
- 2135 Pei, S.W., Gao, X., Chen, F.Y., Feng, X.W., Dennell, R., 2010. Lithic assemblage
2136 from the Jingshuiwan Paleolithic site of the early Late Pleistocene in the Three
2137 Gorges, China. *Quaternary International* **211**, 66–74.
- 2138 Pei, S., Gao, X., Wu, X., Li, X., Bae, C.J., 2013. Middle to Late Pleistocene hominin
2139 occupation in the Three Gorges region, South China. *Quaternary International*
2140 **295**, 237–252.
- 2141 Penkman, K.E.H., 2007. Amino Acid Dating. In: White, T.S., Bridgland, D.R.,
2142 Howard, A.J., White, M.J. (Eds), *The Quaternary of the Trent Valley and*
2143 *Adjoining Areas: Field Guide*. Quaternary Research Association, London, 58–61.
- 2144 Penkman, K.E.H., Preece, R.C., Keen, D.H., Maddy, D., Schreve, D.C., Collins,
2145 M.J., 2007. Testing the aminostratigraphy of fluvial archives: the evidence from
2146 intra-crystalline proteins within freshwater shells. *Quaternary Science Reviews*
2147 **26**, 2958–2969.
- 2148 Penkman, K.E.H., Preece, R.C., Bridgland, D.R., Keen, D.H., Meijer, T., Parfitt, S.A.,
2149 White, T.S., Collins, M.J., 2011. A chronological framework for the British
2150 Quaternary based on Bithynia opercula. *Nature* **476**, 446–449.

- 2151 Penkman, K.E.H., Preece, R.C., Bridgland, D.R., Keen, D.H., Meijer, T., Parfitt, S.A.,
 2152 White, T.S., Collins, M.J., 2013. An aminostratigraphy for the British Quaternary
 2153 based on *Bithynia opercula*. *Quaternary Science Reviews* **61**, 111–134.
- 2154 Pentecost, A., 1995. The Quaternary travertine deposits of Europe and Asia Minor.
 2155 *Quaternary Science Reviews* **14**, 1005–1028.
- 2156 Pereira, J.C., Lopes, R.P., Kerber, L., 2012. New remains of late Pleistocene
 2157 mammals from the Chuí Creek, Southern Brazil. *Revista Brasileira de*
 2158 *Paleontologia* **15**, 228–239.
- 2159 Peretto, C., Arnaud, J., Moggi-Cecchi, J., Manzi, G., Nomade, S., Pereira, A.,
 2160 Falguères, C., Bahain, J.-J., Grimaud-Hervé, D., Berto, C., Sala, B., Lembo, G.,
 2161 Muttilli, B., Gallotti, R., Thun Hohenstein, U., Vaccaro, C., Coltorti, M.,
 2162 Arzarello, M. 2015. A Human Deciduous Tooth and New ⁴⁰Ar/³⁹Ar Dating
 2163 Results from the Middle Pleistocene Archaeological Site of Isernia La Pineta,
 2164 Southern Italy. *PLOS One*, <http://dx.doi.org/10.1371/journal.pone.0140091>.
- 2165 Petraglia, M.D., 2011. Trailblazers across Arabia. *Nature* **470**, 50–51.
- 2166 Petronio, C., Bellucci, L., Martinetto, E., Pandolfi, L., Salari, L., 2011.
 2167 Biochronology and palaeoenvironmental changes from the Middle Pliocene to the
 2168 Late Pleistocene in Central Italy. *Geodiversitas* **33**, 485–517.
- 2169 Pinto Llona, A.C., Andrews, P., 1996. Amphibian taphonomy from cave deposits in
 2170 England. *Comunicación de l Il Reunión de Tafonimía y Fosilización*, 1996,
 2171 327–330.
- 2172 Pinto Llona, A.C., Andrews, P., 1999. Amphibian taphonomy and its application to
 2173 the fossil record of Dolina (Middle Pleistocene, Atapuerca, Spain).
 2174 *Palaeogeography, Palaeoclimatology, Palaeoecology* **149**, 411–429.
- 2175 Porch, N., Kershaw, A.P., 2010. Comparative AMS ¹⁴C dating of plant macrofossils,
 2176 beetles and pollen preparations from two late Pleistocene sites in southeastern
 2177 Australia. In: Haberle, S., Stevenson, J., Prebble, M. (Eds), *Altered Ecologies:*
 2178 *Fire, Climate and Human Influence on Terrestrial Landscapes*. ANU E Press,
 2179 Canberra, 395–404.
- 2180 Powers, R., Ramirez, L., Redmond, C., Elberg, E., 1966. Geology of the Arabian
 2181 Peninsula. Sedimentary Geology of Saudi Arabia Geology of Saudi Arabia.
 2182 USGS Professional paper 560-D. Washington.
- 2183 Prado, J.L., Alberdi, M.T., 2010. Quaternary mammalian faunas of the Pampean
 2184 Region. *Quaternary International* **212**, 176–186.
- 2185 Preece, R.C., 2001. Molluscan evidence for differentiation of interglacials within the
 2186 “cromerian complex”. *Quaternary Science Reviews* **20**, 1643–1656.
- 2187 Preece, R.C., Day, S.P., 1994. Comparison of post-glacial molluscan and vegetational
 2188 successions from a radiocarbon-dated tufa sequence in Oxfordshire. *Journal of*
 2189 *Biogeography* **21**, 463–478.
- 2190 Preece, R.C., Parfitt, S.A., Bridgland, D.R., Lewis, S.G., Rowe, P.J., Atkinson, T.C.,
 2191 Candy, I., Debenham, N.C., Penkman, K.E.H., Rhodes, E.J. and Schwenninger,
 2192 J.L., 2007. Terrestrial environments during MIS 11: evidence from the
 2193 Palaeolithic site at West Stow, Suffolk, UK. *Quaternary Science Reviews* **26**,
 2194 1236–1300.

- 2195 Preece, R.C., Parfitt, S.A., Coope, G.R., Penkman, K.E., Pöhl, P., Whittaker, J.E.,
2196 2009. Biostratigraphic and aminostratigraphic constraints on the age of the
2197 Middle Pleistocene glacial succession in north Norfolk, UK. *Journal of*
2198 *Quaternary Science* **24**, 557–580.
- 2199 Prescott, G.W., Williams, D.R., Balmford, A., Green, R.E., Manica, A., 2012.
2200 Quantitative global analysis of the role of climate and people in explaining late
2201 Quaternary megafaunal extinctions. *Proceedings of the National Academy of*
2202 *Sciences* **109**, 4527–4531.
- 2203 Price, G.J. and Sobbe, I.H., 2005. Pleistocene palaeoecology and environmental
2204 change on the Darling Downs, southeastern Queensland, Australia. *Memoirs of*
2205 *the Queensland Museum* **51**, 171–201.
- 2206 Price, G.J., Webb, G.E., Zhao, J.-X., Feng, Y.-X., Murray, A.S., Cooke, B.N.,
2207 Hocknull, S.A., Sobbe, I.H., 2011. Dating megafaunal extinction on the Pleisto-
2208 cene Darling Downs, eastern Australia: the promise and pitfalls of dating as a test
2209 of extinction hypotheses. *Quaternary Science Reviews* **30**, 899–914.
- 2210 Price, G.J., Louys, J., Cramb, J., Feng, Y.X., Zhao, J.X., Hocknull, S.A., Webb, G.E.,
2211 Nguyen, A.D. and Joannes-Boyau, R., 2015. Temporal overlap of humans and
2212 giant lizards (Varanidae; Squamata) in Pleistocene Australia. *Quaternary Science*
2213 *Reviews* **125**, 98–105.
- 2214 Prideaux, G.J., Roberts, R.G., Megirian, D., Westaway, K.E., Hellstrom, J.C. and
2215 Olley, J.M., 2007. Mammalian responses to Pleistocene climate change in
2216 southeastern Australia. *Geology* **35**, 33–36.
- 2217 Prideaux, G.J., Gully, G.A., Couzens, A.M., Ayliffe, L.K., Jankowski, N.R., Jacobs,
2218 Z., Roberts, R.G., Hellstrom, J.C., Gagan, M.K., Hatcher, L.M., 2010. Timing
2219 and dynamics of Late Pleistocene mammal extinctions in southwestern Australia.
2220 *Proceedings of the National Academy of Sciences* **107**, 22157–22162.
- 2221 Puisségur, J.J., 1976. Mollusques continentaux quaternaires de Bourgogne. Mémoires
2222 géologiques de l'Université de Dijon, Paris.
- 2223 Puchkov, V., Danukalova, G., 2009. The Late Pliocene and Pleistocene history of the
2224 Southern Urals Region in the light of neotectonic data. *Quaternary International*
2225 **201**, 4–12.
- 2226 Ramos, A., Cunha, P.P., Cunha, L., Gomes, A., Lopes, F.C., Buylaert, J.-P., Murray,
2227 A.S., 2012. The River Mondego terraces at the Figueira da Foz coastal area
2228 (western central Portugal): geomorphological and sedimentological
2229 characterization of a terrace staircase affected by differential uplift and glacio-
2230 eustasy. *Geomorphology* **165-166**, 107–123.
- 2231 Read, K.E., 1997. Early hominid evolution and ecological change through the African
2232 Plio-Pleistocene. *Journal of Human Evolution* **32**, 289–322.
- 2233 Reguero, M.A., Candela, A.M., Alonso, R.N., 2007. Biochronology and
2234 biostratigraphy of the Uquia Formation (Pliocene–early Pleistocene, NW
2235 Argentina) and its significance in the Great American Biotic Interchange. *Journal*
2236 *of South American Earth Sciences* **23**, 1–16.
- 2237 Rixhon G., Briant R.M., Cordier S., Duval M., Jones A., Scholz A., 2017 (this issue).
2238 Setting the pace of river landscape evolution at different timescales during the

- 2239 Quaternary: recent developments in numerical dating methods. *Quaternary*
2240 *Science Reviews*.
- 2241 Roe, H.M., 2001. The late Middle Pleistocene biostratigraphy of the Thames Valley,
2242 England: new data from eastern Essex. *Quaternary Science Reviews* **20**, 1603–
2243 1619.
- 2244 Roe, H.M. & Preece, R.C., 1995. A new discovery of the Pleistocene ‘Rhenish’ fauna
2245 in Essex. *Journal of Conchology* **35**, 272–273.
- 2246 Roe, H.M. and Preece, R.C., 2011. Incised palaeo-channels of the late Middle
2247 Pleistocene Thames: age, origins and implications for fluvial palaeogeography
2248 and sea-level reconstruction in the southern North Sea basin. *Quaternary Science*
2249 *Reviews* **30**, 2498–2519.
- 2250 Roe, H.M., Coope, G.R., Devoy, R.J., Harrison, C.J., Penkman, K.E., Preece, R.C.
2251 and Schreve, D.C., 2009. Differentiation of MIS 9 and MIS 11 in the continental
2252 record: vegetational, faunal, aminostratigraphic and sea-level evidence from
2253 coastal sites in Essex, UK. *Quaternary Science Reviews* **28**, 2342–2373.
- 2254 Roe, H.M., Penkman, K.E., Preece, R.C., Briant, R.M. and Wenban-Smith, F.F.,
2255 2011. Evolution of the Thames estuary during MIS 9: insights from the
2256 Shoeburyness area, Essex. *Proceedings of the Geologists' Association* **122**, 397–
2257 418.
- 2258 Rose, J.I., Usik, V.I., Marks, A.E., Hilbert, Y.H., Galletti, C.S., Parton, A., Geiling,
2259 J.M., Černý, V., Morley, M.W., Roberts, R.G., 2011. The Nubian Complex of
2260 Dhofar, Oman: An African Middle Stone Age Industry in Southern Arabia. *PLoS*
2261 *One* **6**, e28239
- 2262 Rosenberg, T.M., Preusser, F., Fleitmann, D., Schwalb, A., Penkman, K., Schmid,
2263 T.W., Al-Shanti, M. a., Kadi, K., Matter, A., 2011. Humid periods in southern
2264 Arabia: windows of opportunity for modern human dispersal. *Geology* **39**, 1115–
2265 1118
- 2266 Rosenberg, T.M., Preusser, F., Risberg, J., Pliikk, A., Kadi, K. a., Matter, A.,
2267 Fleitmann, D., 2013. Middle and Late Pleistocene humid periods recorded in
2268 palaeolake deposits of the Nafud desert, Saudi Arabia. *Quaternary Science*
2269 *Reviews* **70**, 109–123.
- 2270 Rousseau, D.D., Puisségur, J.J., Lécalle, F., 1992. West-European terrestrial molluscs
2271 assemblages of isotopic stage 11 (Middle Pleistocene): climatic implications.
2272 *Palaeogeography, Palaeoclimatology, Palaeoecology* **92**, 15–29.
- 2273 Rule, S., Brook, B.W., Haberle, S.G., Turney, C.S., Kershaw, A.P. and Johnson, C.N.,
2274 2012. The aftermath of megafaunal extinction: ecosystem transformation in
2275 Pleistocene Australia. *Science* **335**, 1483–1486.
- 2276 Sala, B. and Masini, F., 2007. Late Pliocene and Pleistocene small mammal
2277 chronology in the Italian peninsula. *Quaternary International* **160**, 4–16.
- 2278 Sakaguchi, S., Bowman, D.M.J.S., Prior, L.D., Crisp, M.D., Linde, C.C., Tsumura,
2279 Y., Isagi, Y., 2013. Climate, not Aboriginal landscape burning, controlled the
2280 historical demography and distribution of fire-sensitive conifer populations
2281 across Australia. *Proceedings of the Royal Society of London B: Biological*
2282 *Sciences* **280**, 20132182.

- 2283 Sancho, C., Arenas, C., Vázquez-Urbez, M., Pardo, G., Lozano, M.V., Peña-Monné,
2284 J.L., Hellstrom, J., Ortiz, J.E., Osácar, M.C., Auqué, L. and Torres, T., 2015.
2285 Climatic implications of the Quaternary fluvial tufa record in the NE Iberian
2286 Peninsula over the last 500ka. *Quaternary Research* **84**, 398–414.
- 2287 Sancho, C., Calle, M., Peña-Monné, J.L., Duval, M., Oliva-Urcia, B., Pueyo, E.L.,
2288 Benito, G. and Moreno, A., 2016. Dating the Earliest Pleistocene alluvial terrace
2289 of the Alcanadre River (Ebro Basin, NE Spain): Insights into the landscape
2290 evolution and involved processes. *Quaternary International* **407**, 86–95.
- 2291 dom, C., Faurby, S., Sandel, B., Svenning, J.C., 2014. Global late Quaternary
2292 megafauna extinctions linked to humans, not climate change. *Proceedings of the*
2293 *Royal Society B* **281**, 20133254.
- 2294 Santisteban, J.I., Schulte, L., 2007. Fluvial networks of the Iberian Peninsula: a
2295 chronological framework. *Quaternary Science Reviews* **26**, 2738–2757.
- 2296 Sardella, R., Angelote, C., Barisone, G., Bedetti, C., Di Canzio, E., Marcolini, F.,
2297 Squazzini, E., 2003. Italian Plio-Pleistocene mammal biochronology and
2298 correlations with marine sequences: three case studies *II*. *Quaternario* **16**, 15–20.
- 2299 Sardella, R., Bedetti, C., Iurino, D.A., Pavia, M., Bellucci, L., 2015. Comment on
2300 "Reassessing the sedimentary deposits and vertebrate assemblages from Ponte
2301 Galeria area (Rome, central Italy): an archive for the Middle Pleistocene faunas
2302 of Europe" by F. Marra, L. Pandolfi, C. Petronio, G. Di Stefano, M. Gaeta and L.
2303 Salari [*Earth-Science Reviews* 139 (2014), 104–122]. *Earth-Science Reviews* **147**,
2304 201–203.
- 2305 Scerri, E.M.L., Breeze, P.S., Parton, A., Groucutt, H.S., White, T.S., Stimpson, C.,
2306 Clark-balzan, L., Jennings, R., Alsharekh, A., Petraglia, M.D., 2015. Middle to
2307 Late Pleistocene human habitation in the western Nefud Desert, Saudi Arabia.
2308 *Quaternary International* **382**, 200–214.
- 2309 Schreve, D.C., 2001a. Differentiation of the British late Middle Pleistocene
2310 interglacials: the evidence from mammalian biostratigraphy. *Quaternary Science*
2311 *Reviews* **20**, 1693–1705.
- 2312 Schreve, D.C., 2001b. Mammalian evidence from fluvial sequences for complex
2313 environmental change at the oxygen isotope substage level. *Quaternary*
2314 *International* **79**, 65–74.
- 2315 Schreve, D.C., 2009. A new record of Pleistocene hippopotamus from River Severn
2316 terrace deposits, Gloucester, UK - palaeoenvironmental setting and
2317 stratigraphical significance. *Proceedings of the Geologists' Association* **120**, 58–
2318 64.
- 2319 Schreve, D.C., 2012. The Reinsdorf Interglacial (Schöningen II) mammalian
2320 assemblage in its European context. In: Behre, K.-E. (Ed.), *Die chronologische*
2321 *Einordnung der paläolithischen Fundstellen von Schöningen*. *The chronological*
2322 *setting of the Palaeolithic sites of Schöningen*. Forschungen zur Urgeschichte im
2323 Tagebau von Schöningen 1. RGZM, Mainz, 129–142.
- 2324 Schreve, D.C., Bridgland, D.R., 2002. Correlation of English and German Middle
2325 Pleistocene fluvial sequences based on mammalian biostratigraphy. *Netherlands*
2326 *Journal of Geosciences* **81**, 357–373.

- 2327 Schreve, D.C., Candy, I., 2010. Interglacial climates: Advances in our understanding
2328 of warm climate episodes. *Progress in Physical Geography* **34**, 845–856.
- 2329 Schreve, D.C., Keen, D.H., Limondin-Lozouet, N., Auguste, P., Santisteban, J.I.,
2330 Ubilla, M., Matoshko, A., Bridgland, D.R., Westaway, R., 2007. Progress in
2331 faunal correlation of Late Cenozoic fluvial sequences 2000–4: the report of the
2332 IGCP 449 biostratigraphy subgroup. *Quaternary Science Reviews* **26**, 2970–2995.
- 2333 Schreve, D.C., Howard, A.J. Currant, A.P., Brooks, S., Buteux, S., Coope, G.R.,
2334 Crocker, B., Field, M., Greenwood, M., Greig, J., Toms, P., 2013. A Middle
2335 Devensian woolly rhinoceros (*Coelodonta antiquitatis*) from Whitemoor Haye
2336 Quarry, Staffordshire (UK): palaeoenvironmental context and significance.
2337 *Journal of Quaternary Science* **28**, 118–130.
- 2338 Scott, E., 2010. Extinctions, scenarios, and assumptions: changes in latest Pleistocene
2339 large herbivore abundance and distribution in western North America.
2340 *Quaternary International* **217**, 225–239.
- 2341 Scott, B., Ashton, N., Penkman, K.E., Preece, R.C., White, M., 2010. The position
2342 and context of Middle Palaeolithic industries from the Ebbsfleet Valley, Kent,
2343 UK. *Journal of Quaternary Science* **25**, 931–944.
- 2344 Sesé, C., Soto, E. and Pérez González, A., 2000. Mamíferos de las terrazas del valle
2345 del Tajo: primeros datos de micromamíferos del Pleistoceno en Toledo (España
2346 central). *Geogaceta* **28**, 133–136.
- 2347 Seyrek, A., Demir, T., Westaway, R., Guillou, J., Scaillet, S., White, T.S., Bridgland,
2348 D.R., 2014a. The kinematics of central-southern Turkey and northwest Syria
2349 revisited. *Tectonophysics* **618**, 35–66.
- 2350 Seyrek, A., Demir, T., Westaway, R., Guillou, J., Scaillet, S., White, T.S., Bridgland,
2351 D.R., 2014b. Corrigendum to "The kinematics of central-southern Turkey and
2352 northwest Syria revisited." [*Tectonophysics* 618, 35–66]. *Tectonophysics* **630**,
2353 319–320.
- 2354 Shiyake, S., 2014. Applying the Mutual Climatic Range method to the beetle
2355 assemblages in Japan using accurate data of climate and distribution of modern
2356 species. *Quaternary International* **341**, 267–271.
- 2357 Sier, M.J., Parés, J.M., Antoine, P., Loch, J.L., Dekkers, M.J., Limondin-Lozouet, N.
2358 and Roebroeks, W., 2015. Evidence for the Blake Event recorded at the Eemian
2359 archaeological site of Caours, France. *Quaternary International* **357**, 149–157.
- 2360 Sierralta, M., Kele, S., Melcher, F., Hambach, U., Reinders, J., van Geldern, R.,
2361 Frechen, M., 2010. Uranium-series dating of travertine from Süttő: implications
2362 for reconstruction of environmental change in Hungary. *Quaternary International*
2363 **222**, 178–193.
- 2364 Sitzia, L., Bertran, P., Boulogne, S., Brenet, M., Crassard, R., Delagnes, A., Frouin,
2365 M., Hatté, C., Jaubert, J., Khalidi, L., Messenger, E., Mercier, N., Meunier, A.,
2366 Peigné, S., Queffelec, A., Tribolo, C., Macchiarelli, R., 2012. The
2367 Paleoenvironment and Lithic Taphonomy of Shi'bat Dihya 1, a Middle
2368 Paleolithic Site in Wadi Surdud, Yemen. *Geoarchaeology* **27**, 471–491.
- 2369 Soibelzon, E., Prevosti, F.J., Bidegain, J.C., Rico, Y., Verzi, D.H. and Tonni, E.P.,
2370 2009. Correlation of late Cenozoic sequences of southeastern Buenos Aires


- 2371 province: Biostratigraphy and magnetostratigraphy. *Quaternary International*
2372 **210**, 51–56.
- 2373 Stimpson, C.M., Breeze, P., Clark-Balzan, L., Groucutt, H.S., Jennings, R.P., Parton,
2374 A. Scerri, E., White, T.S., Petraglia, M.D., 2015. Stratified Pleistocene
2375 vertebrates with a new record of *Panthera gombaszogensis* (Kretzoi, 1938) from
2376 northern Saudi Arabia. *Quaternary International* **382**, 168–180.
- 2377 Stoetzel, E., Denys, C., Bailon, S., El Hajraoui, M.A., Nespoulet, R., 2012.
2378 Taphonomic analysis of amphibian and squamate remains from El Harhoura 2
2379 (Rabat-Témara, Morocco): contributions to palaeoecological and archaeological
2380 interpretations. *International Journal of Osteoarchaeology* **22**, 616–635.
- 2381 Stuart, A.J., 2015. Late Quaternary megafaunal extinctions on the continents: a short
2382 review. *Geological Journal* **50**, 338–363.
- 2383 Sutcliffe, A.J. (1964). The mammalian fauna. In: Ovey, C.D. (Ed.), *The Swanscombe*
2384 *Skull. A Survey of Research on a Pleistocene Site*. Royal Anthropological
2385 Institute of Great Britain and Ireland, London, Occasional Paper No 20, 85–112.
- 2386 Tchernov, E., 1981. The biostratigraphy of the Middle East. *Préhistoire du Levant*,
2387 Éditions du CNRS, Paris, 67-97.
- 2388 Tchernov, E., 1994. New comments on the biostratigraphy of the Middle and Upper
2389 Pleistocene of the southern Levant. In: Bar-Yosef, O., Kra, R.S. (Eds), *Late*
2390 *Quaternary Chronology and Paleoclimates of the Eastern Mediterranean*.
2391 American School of Prehistoric Research, Tucson, Arizona, 333–350.
- 2392 Thomas, G.N., 2001. Late Middle Pleistocene pollen biostratigraphy in Britain:
2393 pitfalls and possibilities in the separation of interglacial sequences. *Quaternary*
2394 *Science Reviews* **20**, 1621–1630.
- 2395 Thomas, H., Geraads, D., Vaslet, D., Memesh, A., Billiou, D., Bocherens, H.,
2396 Dobigny, G., Eisenmann, V., Gayet, M., Lapparent de Broin, F. de., Petter, G.,
2397 Halawani, M., 1998. First Pleistocene faunas from the Arabian Peninsula: an
2398 Nafud desert, Saudi Arabia. *Comptes Rendus de l'Académie des Sciences* **326**,
2399 145–152.
- 2400 Tonni, E.P., Soibelzon, E., Cione, A.L., Carlinin, A.A., Scillato-Yané, G.J., Zurita,
2401 A.E., Paredes Ríos, F., 2009. Preliminar correlation of the Pleistocene sequences
2402 of the Tarija valley (Bolivia) with the Pampean chronological standard.
2403 *Quaternary International* **210**, 57–65.
- 2404 Ubilla, M., Martínez, S., 2016. *Geology and Paleontology of the Quaternary of*
2405 *Uruguay*. Springer International Publishing, Switzerland.
- 2406 Ubilla, M., Perea, D., Goso, C., Lorenzo, N., 2004. Late Pleistocene vertebrates from
2407 northern Uruguay: tools for biostratigraphic, climatic and environmental
2408 reconstruction. *Quaternary International* **114**, 129–142.
- 2409 Ubilla, M., Perea, D., Rinderknecht, A., Corona, A., 2009. Pleistocene mammals from
2410 Uruguay: biostratigraphic, biogeographic and environmental connotations. In:
2411 Ribeiro, A.M., Bauermann, S.G., Scherer, C.S. (Eds): *Quaternario de Rio*
2412 *Grande do Sul*. Integrando Conhecimentos Monografias Sociedade Brasileira de
2413 Paleontología, Porto Alegre, 217–230.

- 2414 Ubilla, M., Corona, A., Rinderknecht, A., Perea, D., Verde, M., 2016. Marine Isotope
2415 Stage 3 (MIS 3) and Continental Beds from Northern Uruguay (Sopas
2416 Formation): Paleontology, Chronology, and Climate. In: Gasparini, G.M.,
2417 Rabassa, J., Deschamps, C., Tonni, E.P. (Eds): *Marine Isotope Stage 3 in*
2418 *Southern South America, 60 Ka BP-30 Ka BP*. Springer International Publishing,
2419 Switzerland, 183–205.
- 2420 Urban, B., Bigga, G., 2015. Environmental reconstruction and biostratigraphy of late
2421 Middle Pleistocene lakeshore deposits at Schöningen. *Journal of Human*
2422 *Evolution* **89**, 57–70.
- 2423 Vaks, A., Bar-Matthews, M., Ayalon, A., Matthews, A., Halicz, L., Frumkin, A.,
2424 2007. Desert speleothems reveal climatic window for African exodus of early
2425 modern humans. *Geology* **35**, 831.
- 2426 Vaks, A., Woodhead, J., Bar-Matthews, M., Ayalon, A., Cliff, R. A., Zilberman, T.,
2427 Matthews, A., Frumkin, A., 2013. Pliocene–Pleistocene climate of the northern
2428 margin of Saharan–Arabian Desert recorded in speleothems from the Negev
2429 Desert, Israel. *Earth and Planetary Science Letters* **368**, 88–100.
- 2430 Vallverdú, J., Saladié, P., Rosas, A., Huguet, R., Cáceres, I., Mosquera, M., Garcia-
2431 Tabernero, A., Estalrich, A., Lozano-Fernández, I., Pineda-Alcalá, A.,
2432 Carrancho, A., Villalain, J.J., Bourlès, D., Braucher, R., Lebatard, A., Vilalta, J.,
2433 Esteban-Nadal, M., Lluç Bennàsar, M., Bastir, M., López-Polín, L., Ollé, A.,
2434 Vergés, J.M., Ros-Montoya, S., Martínez-Navarro, B., García, A., Martinell, J.,
2435 Expósito, I., Burjachs, F., Agustí, J., Carbonell, E., 2014. Age and date for early
2436 arrival of the Acheulian in Europe (Barranc de la Boella, la Canonja, Spain).
2437 *PLOS One* **9**, e103634.
- 2438 Vandenberghe, J., 2015. River terraces as a response to climatic forcing: formation
2439 processes, sedimentary characteristics and sites for human occupation.
2440 *Quaternary International* **370**, 3–11.
- 2441 Vanhoorne, R., 2003. A contribution to the palaeontological study of the Middle
2442 Pleistocene deposits at Lo (Belgium). *Quaternaire* **14**, 75–83.
- 2443 Van Kolfschoten, T., Buhrs, E., Verheijen, I., 2015. The larger mammal fauna from
2444 the Lower Paleolithic Schöningen Spear site and its contribution to hominin
2445 subsistence. *Journal of Human Evolution* **89**, 138–153.
- 2446 Vis, G.J., Kasse, C., Vandenberghe, J., 2008. Late Pleistocene and Holocene
2447 palaeogeography of the Lower Tagus Valley (Portugal): effects of relative sea
2448 level, valley morphology and sediment supply. *Quaternary Science Reviews* **27**,
2449 1682–1709.
- 2450 Vis, G.J., Bohncke, S.J., Schneider, H., Kasse, C., Coenraads - Nederveen, S.,
2451 Zuurbier, K. and Rozema, J., 2010. Holocene flooding history of the lower Tagus
2452 valley (Portugal). *Journal of Quaternary Science* **25**, 1222–1238.
- 2453 Viveen, W., Braucher, R., Bourlès, D., Schoorl, J.M., Veldkamp, A., Van Balen, R.T.,
2454 Wallinga, J., Fernandez-Mosquera, D., Vidal-Romani, J.R., Sanjurjo-Sanchez, J.,
2455 2012. A 0.65 Ma chronology and incision rate assessment of the NW Iberian
2456 Miño River terraces based on 10 Be and luminescence dating. *Global and*
2457 *Planetary Change* **94**, 82–100.

- 2458 Viveen, W., Schoorl, J.M., Veldkamp, A., van Balen, R.T., Vidal-Romani, J.R., 2013.
2459 Fluvial terraces of the northwest Iberian lower Miño River. *Journal of Maps* **9**,
2460 513–522.
- 2461 Voinchet, P., Moreno, D., Bahain, J.J., Tissoux, H., Tombret, O., Falguères, C.,
2462 Moncel, M.H., Schreve, D., Candy, I., Antoine, P., Ashton, N., Beamish, M.,
2463 Cliquet, D., Despriée, J., Lewis, S., Limondin-Lozouet, N., Locht, J.L., Parfitt, S.,
2464 Pope, M., 2015. New chronological data (ESR and ESR/U-series) for the earliest
2465 Acheulian sites of north-western Europe. *Journal of Quaternary Science* **30**, 610–
2466 622.
- 2467 Webb, S., 2008. Megafauna demography and Late Quaternary climatic change in
2468 Australia: a predisposition to extinction. *Boreas* **37**, 329–345.
- 2469 Webb, S., 2009. Late Quaternary distribution and biogeography of the southern Lake
2470 Eyre basin (SLEB) megafauna, South Australia. *Boreas* **38**, 25–38.
- 2471 Wenban-Smith, F.F., Allen, P., Bates, M.R., Parfitt, S.A., Preece, R.C., Stewart, J.R.,
2472 Turner, C., Whittaker, J.E., 2006. The Clactonian elephant butchery site at
2473 Southfleet Road, Ebbsfleet, UK. *Journal of Quaternary Science* **21**, 471–483.
- 2474 Wenban-Smith, F.F., 2013. *The Ebbsfleet Elephant. Excavations at Southfleet Road,*
2475 *Swanscombe, in advance of High Speed 1, 2003-4.* Oxford Archaeology
2476 Monograph No. 20, Oxford Archaeology, Oxford.
- 2477 Wenban-Smith, F.F., 2014. Baker's Hole, Ebbsfleet (TL 615742) Areas B and F:
2478 Burchell's Ebbsfleet Channel site (TQ 561240) & Temperate Bed (TQ 174050).
2479 In: Bridgland, D.R., Allen, P., White, T.S. (Eds.) *The Quaternary of the Lower*
2480 *Thames and eastern Essex. Field Guide.* Quaternary Research Association,
2481 London, 189–198.
- 2482 Westaway, K.E., Morwood, J.M., Roberts, R.G., Rokus, A.D., Zhao, J.X., Storm, P.,
2483 Aziz, F., van den Bergh, G., Hadi, P., Jatmicode Vos, J., 2007. Age and
2484 biostratigraphic significance of the Punung Rainforest fauna, East Java,
2485 Indonesia, and implications for Pongo and Homo. *Journal of Human Evolution*
2486 **53**, 709–717.
- 2487 Westaway, R., 2009. Calibration of decomposition of serine to alanine in Bithynia
2488 opercula as a quantitative dating technique for Middle and Late Pleistocene sites
2489 in Britain. *Quaternary Geochronology* **4**, 241–259.
- 2490 Westaway, R., 2017. Isostatic compensation of Quaternary vertical crustal motions:
2491 coupling between uplift of Britain and subsidence beneath the North Sea. *Journal*
2492 *of Quaternary Science* **32**, 169–182.
- 2493 Westaway, R., Bridgland, D.R., Sinha, R., Demir, T., 2009. Fluvial sequences as
2494 evidence for landscape and climatic evolution in the Late Cenozoic: a synthesis
2495 of data from IGCP 518. *Global and Planetary Change* **68**, 237–253.
- 2496 Westaway R., Bridgland D.R., White T.S., Howard, A.J., White, M.J., 2015. The use
2497 of uplift modelling in the reconstruction of drainage development and landscape
2498 evolution in the repeatedly glaciated Trent catchment, English Midlands, UK.
2499 *Proceedings of the Geologists' Association* **126**, 480–521.
- 2500 Westaway, M.C., Olley, J., Grün, R., 2017. At least 17,000 years of coexistence:
2501 Modern humans and megafauna at the Willandra Lakes, South-Eastern
2502 Australia. *Quaternary Science Reviews* **157**, 206–211.

- 2503 White, M., 2016. Lost Landscapes of Palaeolithic Britain. The contribution of projects
2504 funded by the Aggregates Levy Sustainability Fund 2002–2011. Oxford
2505 Archaeology Monograph 26, Oxford Archaeology, Oxford.
- 2506 White, T.S., 2012. *Late Middle Pleistocene non-marine mollusc and ostracod*
2507 *successions and their relevance to Palaeolithic archaeology*. Unpublished Ph.D.
2508 Thesis, University of Cambridge.
- 2509 White, T.S., Bridgland, D.R., Howard, A.J., White, M.J. 2007. *Quaternary of the*
2510 *Trent Valley and Adjoining Regions: Field Guide*. Quaternary Research
2511 Association, London.
- 2512 White, T.S., Bridgland, D.R., Howard, A.J., White, M.J., Westaway, R.W.C., 2010.
2513 Evidence from the Trent terrace archive, Lincolnshire, UK, for lowland glaciation
2514 of Britain during the Middle and Late Pleistocene. *Proceedings of the Geologists'*
2515 *Association* **121**, 141–153.
- 2516 White, T.S., Preece, R.C. and Whittaker, J.E., 2013. Molluscan and ostracod
2517 successions from Dierden's Pit, Swanscombe: insights into the fluvial history,
2518 sea-level record and human occupation of the Hoxnian Thames. *Quaternary*
2519 *Science Reviews* **70**, 73–90.
- 2520 White, T.S., Bridgland, D.R., Westaway, R.W.C., Straw, A., 2017. Evidence for late
2521 Middle Pleistocene glaciation of the British margin of the southern North Sea.
2522 *Journal of Quaternary Science* **32**, 261–275
- 2523 Whittaker, J.E., Horne, D.J., 2009. Pleistocene. In: Whittaker, J.E., Hart, M.B. (Eds)
2524 *Ostracods in British Stratigraphy*. Micropalaeontological Society Special
2525 Publication, Geological Society Publishing House, Bath, 447–468.
- 2526 Woodburne, M.O., 2010. The Great American Biotic Interchange: Dispersals,
2527 Tectonics, Climate, Sea Level and Holding Pens. *Journal of Mammal Evolution*
2528 **17**, 245–264.
- 2529 Woodburne, M.O., 2012. *Late Cretaceous and Cenozoic mammals of North America:*
2530 *biostratigraphy and geochronology*. Columbia University Press.
- 2531 Wymer, J.J., 1999. *The Lower Palaeolithic Occupation of Britain*. Wessex
2532 Archaeology and English Heritage, Salisbury.
- 2533 Yakovlev, A., Danukalova, G., Osipova, E., 2013. Biostratigraphy of the Upper
2534 Pleistocene (Upper Neopleistocene) of the Southern Urals. *Quaternary*
2535 *International* **292**, 150–167.
- 2536 Yang, G., Zhang, X., Tian, M., Ping, Y., Chen, A., Ge, Z., Ni, Z., Yang, Z., 2011.
2537 Geomorphological and sedimentological comparison of fluvial terraces and karst
2538 caves in Zhangjiajie, northwest Hunan, China: an archive of sandstone landform
2539 development. *Environmental Earth Sciences* **64**, 671–683.
- 2540 Yanina, T.A., 2013. Biostratigraphy of the Middle and Upper Pleistocene of the
2541 Caspian region. *Quaternary International* **284**, 85–97.
- 2542 Yonge, D., Hesse, P.P., 2009. Geomorphic environments, drainage breakdown, and
2543 channel and floodplain evolution on the lower Macquarie River, central-western
2544 New South Wales. *Australian Journal of Earth Sciences* **56**, S35–S53.

- 2545 Zagwijn, W.H., 1996. The Cromerian Complex Stage of the Netherlands and
 2546 correlation with other areas in Europe. In: Turner, C. (Ed.) *The Early Middle*
 2547 *Pleistocene in Europe*. Balkema, Rotterdam, 145–172.
- 2548 Zettler, M.L., 2008. Zur Taxonomie und Verbreitung der Gattung *Theodoxus*
 2549 Montfort, 1810 in Deutschland. Darstellung historischer und rezenter Daten
 2550 einschließlich einer Bibliografie. *Mollusca* **26**, 13–72.
- 2551 Zhao, H., Lu, Y., Wang, C., Chen, J., Liu, J., Mao, H., 2010. ReOSL dating of aeolian
 2552 and fluvial sediments from Nihewan Basin, northern China and its environmental
 2553 application. *Quaternary Geochronology* **5**, 159–163.
- 2554 Zurita, A.E., Miño-Boilini, Á.R., Soibelzon, E., Carlini, A.A., Ríos, F.P., 2009a. The
 2555 diversity of Glyptodontidae (Xenarthra, Cingulata) in the Tarija Valley (Bolivia):
 2556 systematic, biostratigraphic and paleobiogeographic aspects of a particular
 2557 assemblage. *Neues Jahrbuch für Geologie und Paläontologie-Abhandlungen* **251**,
 2558 225–237.
- 2559 Zurita, A.E., Carlini, A.A., Scillato-Yané, G.J., 2009b. Paleobiogeography,
 2560 biostratigraphy and systematics of the Hoplophorini (Xenarthra, Glyptodontidae,
 2561 Hoplophorinae) from the Ensenadan Stage (early Pleistocene to early Middle
 2562 Pleistocene). *Quaternary International* **210**, 82–92.
- 2563






Figure
[Click here to download high resolution image](#)







Figure
[Click here to download high resolution image](#)




La Celle,
late Middle Pleistocene (MIS 11)



Caours,
Late Pleistocene (MIS 5e)




Saint Germain le Vasson,
Holocene (MIS 1)



Modern European species ranges:
N: North; W: West; C: Central; E: East; S: South

Figure

[Click here to download high resolution image](#)



[Click here to download high resolution image](#)

Mammal Isotope Stage	Chronology	Glacial/Interglacial Stage	Key Deposits	Large Mammal Assemblages	Elephant phylogeny	Small Mammal Assemblages	Small Mammal Localities	First Appearances of Notable Species	
1	HOLOCENE								
2	LATE PLEISTOCENE	Valda (Dudinka)	Late Valda	TB, bones Trukhtsevsk paleosol TB, bones	SAMAROVITZ	BUNGARAN	Tatara Kudmoo Khokhlova 2 Elisevichi	Dromotrypa angustata	
3			Middle Valda	Bransk, Kuvandrina, Strelshak paleosols			Anapovsk Sungir Troika 2		
4			Early Valda	Lessa Kudlak paleosol Lessa			Selovo Kabani 2 (SR) Gubach	Dromotrypa gubachi	
5		Mikhurin Interglacial	Salyr paleosol	Shuntovin			Shuntovin	Makhuro Chernykh Makhuro 5 Shuntov Elizovo Nemetskaya	Auritus leucurus Musculus (Tartarus submontanus)
6	MIDDLE PLEISTOCENE	Dnieper (Dudinka)	Moscow Stage	TB, bones	CHOKAZARI	CHOKAZARI	Pavlovka-in-Gresh Koltykale Zhukovichi Ripakov Spasivoe Vilgno Strigovo Ochekai (Homo-glacial layer) Yagobche	Dromotrypa angustata	
7			Kursk Interglacial	Kursk paleosol					
8			Dnieper Stage	TB, bones					
9		Warming Cooling	Rumny paleosol Ouchik bones						
10		Kamarka Interglacial	Kamarka paleosol						
11		Cooling	Borzhomsk less						
12		Lutvin Interglacial	Inuhova paleosol						
13		Oka Glaciation	Oka TB, bones						
14		Konakhovka optimum Pudabinsk cooling Slavuta optimum	Vanya soil complex						
15		Don Glaciation	TB, bones						
16	EARLY PLEISTOCENE	Late Vnuka Interglacial	Voronezh paleosol	TRASPOLAN	TRASPOLAN	Cherkash Gunka Chagin Vol. Enencha Rerava Rybnya Golota Olshane Uman Trespol (rich soil) Mihailovka 2 Smolenki Boia	Auritus cantabrigiae		
17						Cooling	Lessa		
18						Middle Vnuka Interglacial	Ternouka paleosol		
19						Cooling	Lessa		
20-28						Early Vnuka Interglacial	Troika paleosol		
21		Cooling	Polotsk less						
22		Baleshevsk Interglacial	Baleshevsk paleosol						
23		Cooling	Tchernensk less						
24		Interglacial	Kiskensk paleosol						
25		Cooling	Lessa						
26	Warming	Khadzhimys paleosol	PSEKUPAN (Late Phase)	Arctostaphyle meridionalis Samanensis	ODESSAN (Late Phase)	Khadzhimys Melniko	Prolegus demoponticus		

Figure
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)





Figure
[Click here to download high resolution image](#)



Highlights

- Research over the last decade into biostratigraphical data recorded in fluvial archives is reviewed
- Consideration of biogeography and palaeoclimatic evidence is also included
- This emphasizes the disparities in research priorities and fossil preservation on a global scale